

Publicazioni di Riccardo Paoletti

1) [The EEE Project: Cosmic rays, multigap resistive plate chambers and high school students.](#)

By M. Abbrescia, S. Aiola, R. Antolini, C. Avanzini, R. Baldini Ferroli, G. Bencivenni, E. Bossini, E. Bressan et al..

[10.1088/1748-0221/7/11/P11011.](#)

JINST 7 (2012) P11011.

2) [Observations of the magnetars 4U 0142+61 and 1E 2259+586 with the MAGIC telescopes.](#)

By MAGIC Collaboration (, et al.).

[arXiv:1211.1173 [Unknown]].

3) [The PMT-block Test Bench Developed in Pisa.](#)

By D. Costanzo, T Del Prete, E. Mazzoni, GF Pagani, R. Paoletti, D. Rizzi.

4) [Jet Finder Library: version 1.0.](#)

By M. Bosman, I.C. Park, M. Cobal, D. Costanzo, S. Lami, R. Paoletti, Georges Azuelos, K. Strahl.

5) [Particle Level Study of W -> qqbar and Jet Algorithms.](#)

By S. Bettelli, A. Bocci, D. Costanzo, S. Kuhlmann, S. Lami, R. Paoletti.

6) [Search for Gravitational Waves from Binary Black Hole Inspiral, Merger and Ringdown in LIGO-Virgo Data from 2009-2010.](#)

By LIGO Scientific and Virgo Collaborations (, et al.).

[arXiv:1209.6533 [gr-qc]].

7) [Einstein@Home all-sky search for periodic gravitational waves in LIGO S5 data.](#)

By The LIGO Scientific and the Virgo Collaboration (J. Aasi et al.).

[arXiv:1207.7176 [gr-qc]].

8) [MAGIC observations of the giant radio galaxy M87 in a low-emission state between 2005 and 2007.](#)

By MAGIC Collaboration (, et al.).

[arXiv:1207.2147 [astro-ph.HE]].

Astron.Astrophys. 544 (2012) A96.

9) [High zenith angle observations of PKS 2155-304 with the MAGIC-I telescope.](#)

By MAGIC Collaboration (J. Aleksic et al.).

[arXiv:1207.1634 [astro-ph.HE]].

10) [Noise monitor tools and their application to Virgo data.](#)

By T. Accadia, F. Acernese, M. Agathos, P. Astone, G. Ballardin, F. Barone, M. Barsuglia, A. Basti et al..

[10.1088/1742-6596/363/1/012024.](#)

J.Phys.Conf.Ser. 363 (2012) 012024.

11) [The NoEMi \(Noise Frequency Event Miner\) framework.](#)

By T. Accadia, F. Acernese, M. Agathos, P. Astone, G. Ballardin, F. Barone, M. Barsuglia, A. Basti et al..

[10.1088/1742-6596/363/1/012037.](#)

J.Phys.Conf.Ser. 363 (2012) 012037.

12) [A First Search for coincident Gravitational Waves and High Energy Neutrinos using LIGO, Virgo and ANTARES data from 2007.](#)

By LIGO Scientific and Virgo Collaborations (S. Adrian-Martinez et al.).

[arXiv:1205.3018 [astro-ph.HE]].

13) [Search for gravitational waves associated with gamma-ray bursts during LIGO science run 6 and Virgo science runs 2 and 3.](#)

By LIGO Scientific Collaboration (J. Abadie et al.).
[arXiv:1205.2216 [astro-ph.HE]].
[10.1088/0004-637X/760/1/12](https://doi.org/10.1088/0004-637X/760/1/12).
Astrophys.J. 760 (2012) 12.

14) [Swift follow-up observations of candidate gravitational-wave transient events.](#)
By LIGO Scientific Collaboration (P.A. Evans et al.).
[arXiv:1205.1124 [astro-ph.HE]].
[10.1088/0067-0049/203/2/28](https://doi.org/10.1088/0067-0049/203/2/28).
Astrophys.J.Suppl. 203 (2012) 28.

15) [Virgo: a laser interferometer to detect gravitational waves.](#)
By VIRGO Collaboration (T. Accadia et al.).
[10.1088/1748-0221/7/03/P03012](https://doi.org/10.1088/1748-0221/7/03/P03012).
JINST 7 (2012) P03012.

16) [The characterization of Virgo data and its impact on gravitational-wave searches.](#)
By VIRGO Collaboration (J. Aasi et al.).
[arXiv:1203.5613 [gr-qc]].
[10.1088/0264-9381/29/15/155002](https://doi.org/10.1088/0264-9381/29/15/155002).
Class.Quant.Grav. 29 (2012) 155002.

17) [Detection of VHE \$\gamma\$ -rays from HESS J0632+057 during the 2011 February X-ray outburst with the MAGIC Telescopes.](#)
By MAGIC Collaboration (, et al.).
[arXiv:1203.2867 [astro-ph.HE]].

18) [Discovery of VHE gamma-rays from the blazar 1ES 1215+303 with the MAGIC Telescopes and simultaneous multi-wavelength observations.](#)
By MAGIC Collaboration (, et al.).
[arXiv:1203.0490 [astro-ph.HE]].
Astron.Astrophys. 544 (2012) A142.

19) [Morphological and spectral properties of the W51 region measured with the MAGIC telescopes.](#)
By MAGIC Collaboration (J. Aleksic et al.).
[arXiv:1201.4074 [astro-ph.HE]].

20) [Discovery of VHE gamma-ray emission from the BL Lac object B3 2247+381 with the MAGIC telescopes.](#)
By MAGIC Collaboration (, et al.).
[arXiv:1201.2634 [astro-ph.HE]].
Astron.Astrophys. 539 (2012) A118.

21) [Detection of very high energy gamma-ray emission from NGC 1275 by the MAGIC telescopes.](#)
By MAGIC Collaboration (J. Aleksic et al.).
[arXiv:1112.3917 [astro-ph.HE]].

22) [Detection of the gamma-ray binary LS I +61 303 in a low flux state at Very High Energy gamma-rays with the MAGIC Telescopes in 2009.](#)
By MAGIC Collaboration (J. Aleksic et al.).
[arXiv:1111.6572 [astro-ph.HE]].
[10.1088/0004-637X/746/1/80](https://doi.org/10.1088/0004-637X/746/1/80).
Astrophys.J. 746 (2012) 80.

23) [The 2010 very high energy gamma-ray flare & 10 years of multi-wavelength observations of M 87.](#)
By H.E.S.S. and VERITAS Collaborations (A. Abramowski et al.).

[arXiv:1111.5341 [astro-ph.CO]].
[10.1088/0004-637X/746/2/151](https://doi.org/10.1088/0004-637X/746/2/151).
Astrophys.J. 746 (2012) 151.

24) [Constraining Cosmic Rays and Magnetic Fields in the Perseus Galaxy Cluster with TeV observations by the MAGIC telescopes.](#)

By MAGIC Collaboration (J. Aleksic et al.).
[arXiv:1111.5544 [astro-ph.HE]].
Astron.Astrophys. 541 (2012) A99.

25) [MAGIC contributions to the 32nd International Cosmic Ray Conference.](#)

By MAGIC Collaboration (J. Aleksic et al.).
[arXiv:1111.0879 [astro-ph.HE]].

26) [Phase-resolved energy spectra of the Crab Pulsar in the range of 50-400 GeV measured with the MAGIC Telescopes.](#)

By MAGIC Collaboration (J. Aleksic et al.).
[arXiv:1109.6124 [astro-ph.HE]].
Astron.Astrophys. 540 (2012) A69.

27) [Observations of the Crab pulsar between 25 GeV and 100 GeV with the MAGIC I telescope.](#)

By MAGIC Collaboration (J. Aleksic et al.).
[arXiv:1108.5391 [astro-ph.HE]].
[10.1088/0004-637X/742/1/43](https://doi.org/10.1088/0004-637X/742/1/43).
Astrophys.J. 742 (2011) 43.

28) [Performance of the MAGIC stereo system obtained with Crab Nebula data.](#)

By MAGIC Collaboration (J. Aleksic et al.).
[arXiv:1108.1477 [astro-ph.IM]].
[10.1016/j.astropartphys.2011.11.007](https://doi.org/10.1016/j.astropartphys.2011.11.007).
Astropart.Phys. 35 (2012) 435-448.

29) [Observation of the February 2011 Forbush decrease by the EEE telescopes.](#)

By M. Abbrescia, S. Aiola, R. Antolini, C. Avanzini, R. Baldini Ferroli, G. Bencivenni, E. Bossini, E. Bressan et al..
[10.1140/epjp/i2011-11061-5](https://doi.org/10.1140/epjp/i2011-11061-5).
Eur.Phys.J.Plus 126 (2011) 61.

30) [Fermi large area telescope observations of Markarian 421: The missing piece of its spectral energy distribution.](#)

By MAGIC Collaboration (A.A. Abdo et al.).
[10.1088/0004-637X/736/2/131](https://doi.org/10.1088/0004-637X/736/2/131).
Astrophys.J. 736 (2011) 131.

31) [Mrk 421 active state in 2008: the MAGIC view, simultaneous multi-wavelength observations and SSC model constrained.](#)

By J. Aleksic, E.A. Alvarez, L.A. Antonelli, P. Antoranz, M. Asensio, M. Backes, J.A. Barrio, D. Bastieri et al..
[arXiv:1106.1589 [astro-ph.HE]].
Astron.Astrophys. 542 (2012) A100.

32) [A search for Very High Energy gamma-ray emission from Scorpius X-1 with the MAGIC telescopes.](#)

By MAGIC Collaboration (J. Aleksic et al.).
[arXiv:1103.5677 [astro-ph.HE]].
Astrophys.J. 735 (2011) L5.

- 33) [Searches for Dark Matter annihilation signatures in the Segue 1 satellite galaxy with the MAGIC-I telescope.](#)
By MAGIC Collaboration (J. Aleksic et al.).
[arXiv:1103.0477 [astro-ph.HE]].
[10.1088/1475-7516/2011/06/035](https://doi.org/10.1088/1475-7516/2011/06/035).
JCAP 1106 (2011) 035.
- 34) [MAGIC discovery of VHE Emission from the FSRQ PKS 1222+21.](#)
By MAGIC Collaboration (J. Aleksic et al.).
[arXiv:1101.4645 [astro-ph.HE]].
Astrophys.J. 730 (2011) L8.
- 35) [PG 1553+113: five years of observations with MAGIC.](#)
By MAGIC Collaboration (J. Aleksic et al.).
[arXiv:1101.2764 [astro-ph.CO]].
[10.1088/0004-637X/748/1/46](https://doi.org/10.1088/0004-637X/748/1/46).
Astrophys.J. 748 (2012) 46.
- 36) [MAGIC observations and multiwavelength properties of the quasar 3C279 in 2007 and 2009.](#)
By MAGIC Collaboration (J. Aleksic et al.).
[arXiv:1101.2522 [astro-ph.CO]].
- 37) [Spectral Energy Distribution of Markarian 501: Quiescent State vs. Extreme Outburst.](#)
By VERITAS and MAGIC Collaborations (V.A. Acciari et al.).
[arXiv:1012.2200 [astro-ph.HE]].
[10.1088/0004-637X/729/1/2](https://doi.org/10.1088/0004-637X/729/1/2).
Astrophys.J. 729 (2011) 2.
- 38) [Insights Into the High-Energy Gamma-ray Emission of Markarian 501 from Extensive Multifrequency Observations in the Fermi Era.](#)
By LAT and MAGIC and VERITAS Collaborations (A.A. Abdo et al.).
[arXiv:1011.5260 [astro-ph.HE]].
[10.1088/0004-637X/727/2/129](https://doi.org/10.1088/0004-637X/727/2/129).
Astrophys.J. 727 (2011) 129.
- 39) [Prospects for Diffractive and Forward Physics at the LHC.](#)
By The CMS and TOTEM diffractive and forward physics working group Collaboration (M. Albrow et al.).
- 40) [Observations of the Blazar 3C 66A with the MAGIC Telescopes in Stereoscopic Mode.](#)
By MAGIC Collaboration (J. Aleksic et al.).
[arXiv:1010.0550 [astro-ph.HE]].
[10.1088/0004-637X/726/2/58](https://doi.org/10.1088/0004-637X/726/2/58).
Astrophys.J. 726 (2011) 58.
- 41) [Detection of very high energy gamma-ray emission from the Perseus cluster head-tail galaxy IC 310 by the MAGIC telescopes.](#)
By MAGIC Collaboration (J. Aleksic et al.).
[arXiv:1009.2155 [astro-ph.HE]].
Submitted to: Astrophys.J..
- 42) [Design concepts for the Cherenkov Telescope Array CTA: An advanced facility for ground-based high-energy gamma-ray astronomy.](#)
By CTA Consortium Collaboration (M. Actis et al.).
[arXiv:1008.3703 [astro-ph.IM]].
[10.1007/s10686-011-9247-0](https://doi.org/10.1007/s10686-011-9247-0).
Exper.Astron. 32 (2011) 193-316.

- 43) [MAGIC Upper Limits for two Milagro-detected, Bright Fermi Sources in the Region of SNR G65.1+0.6.](#)
By J. Aleksic, L.A. Antonelli, P. Antoranz, M. Backes, J.A. Barrio, D. Bastieri, J. Becerra Gonzalez, W. Bednarek et al..
[arXiv:1007.3359 [astro-ph.HE]].
[10.1088/0004-637X/725/2/1629.](#)
Astrophys.J. 725 (2010) 1629-1632.
- 44) [Magic constraints on Gamma-ray emission from Cygnus X-3.](#)
By MAGIC Collaboration (J. Aleksic et al.).
[arXiv:1005.0740 [astro-ph.HE]].
[10.1088/0004-637X/721/1/843.](#)
Astrophys.J. 721 (2010) 843-855.
- 45) [MAGIC observation of the GRB080430 afterglow.](#)
By MAGIC Collaboration (J. Aleksic et al.).
[arXiv:1004.3665 [astro-ph.HE]].
- 46) [Search for an extended VHE gamma-ray emission from Mrk 421 and Mrk 501 with the MAGIC Telescope.](#)
By J. Aleksic, L.A. Antonelli, P. Antoranz, M. Backes, C. Baixeras, J.A. Barrio, D. Bastieri, J. Becerra Gonzalez et al..
[arXiv:1004.1093 [astro-ph.HE]].
- 47) [Gamma-ray excess from a stacked sample of high- and intermediate-frequency peaked blazars observed with the MAGIC telescope.](#)
By MAGIC Collaboration (J. Aleksic et al.).
[arXiv:1002.2951 [astro-ph.HE]].
[10.1088/0004-637X/729/2/115.](#)
Astrophys.J. 729 (2011) 115.
- 48) [MAGIC TeV Gamma-Ray Observations of Markarian 421 during Multiwavelength Campaigns in 2006.](#)
By MAGIC Collaboration (J. Aleksic et al.).
[arXiv:1001.1291 [astro-ph.CO]].
Astron.Astrophys. 519 (2010) A32.
- 49) [Simultaneous multi-frequency observation of the unknown redshift blazar PG 1553+113 in March-April 2008.](#)
By MAGIC Collaboration (J. Aleksic et al.).
[arXiv:0911.1088 [astro-ph.HE]].
Astron.Astrophys. 515 (2010) A76.
- 50) [Correlated X-ray and Very High Energy emission in the gamma-ray binary LS I +61 303.](#)
By MAGIC Collaboration (H. Anderhub et al.).
[arXiv:0910.4381 [astro-ph.HE]].
[10.1088/0004-637X/706/1/L27.](#)
Astrophys.J. 706 (2009) L27-L32.
- 51) [Simultaneous Multiwavelength observation of Mkn 501 in a low state in 2006.](#)
By MAGIC Collaboration (H. Anderhub et al.).
[arXiv:0910.2093 [astro-ph.HE]].
[10.1088/0004-637X/705/2/1624.](#)
Astrophys.J. 705 (2009) 1624-1631.
- 52) [MAGIC Gamma-Ray Telescope Observation of the Perseus Cluster of Galaxies: Implications for](#)

[Cosmic Rays, Dark Matter and NGC 1275.](#)

By MAGIC Collaboration (J. Aleksic et al.).

[arXiv:0909.3267 [astro-ph.HE]].

[10.1088/0004-637X/710/1/634.](#)

Astrophys.J. 710 (2010) 634-647.

53) [Radio Imaging of the Very-High-Energy Gamma-Ray Emission Region in the Central Engine of a Radio Galaxy.](#)

By VERTIAS, HESS and MAGIC Collaboration (V.A. Acciari et al.).

[arXiv:0908.0511 [astro-ph.HE]].

[10.1126/science.1175406.](#)

Science 325 (2009) 444-448.

54) [MAGIC Collaboration: Contributions to the 31st International Cosmic Ray Conference \(ICRC 2009\).](#)

By MAGIC Collaboration (: H. Anderhub et al.).

[arXiv:0907.0843 [astro-ph.HE]].

55) [The readout system of the MAGIC-II Cherenkov Telescope.](#)

By MAGIC Collaboration (D. Tesaro et al.).

[arXiv:0907.0466 [astro-ph.IM]].

56) [Suzaku and Multi-wavelength Observations of OJ 287 during the Periodic Optical Outburst in 2007.](#)

By KANATA and MAGIC Collaboration (Hiromi Seta et al.).

[arXiv:0906.0234 [astro-ph.CO]].

57) [Search for VHE \$\gamma\$ -ray emission from the globular cluster M13 with the MAGIC telescope.](#)

By MAGIC Collaboration (Hans Anderhub et al.).

[arXiv:0905.2427 [astro-ph.HE]].

[10.1088/0004-637X/702/1/266.](#)

Astrophys.J. 702 (2009) 266-269.

58) [High performances and low cost front-end electronics for the Cherenkov Telescope Array.](#)

By CTA Collaboration (P. Vincent et al.).

[10.1063/1.3076820.](#)

AIP Conf.Proc. 1085 (2009) 890-893.

59) [MAGIC Observations of PG 1553+113 during a Multiwavelength Campaign in July 2006.](#)

By MAGIC Collaboration (J. Albert et al.).

[arXiv:0812.3037 [astro-ph]].

60) [The June 2008 flare of Markarian 421 from optical to TeV energies.](#)

By Veritas Collaboration (V. Vittorini et al.).

[arXiv:0812.1500 [astro-ph]].

[10.1088/0004-637X/691/1/L13.](#)

Astrophys.J. 691 (2009) L13-L19.

61) [MAGIC upper limits to the VHE gamma-ray flux of 3C454.3 in high emission state.](#)

By MAGIC Collaboration (Hans Anderhub et al.).

[arXiv:0811.1680 [astro-ph]].

62) [Discovery of a very high energy gamma-ray signal from the 3C 66A/B region.](#)

By MAGIC Collaboration (E. Aliu et al.).

[arXiv:0810.4712 [astro-ph]].

63) [Improving the performance of the single-dish Cherenkov telescope MAGIC through the use of signal timing.](#)

By MAGIC Collaboration (E. Aliu et al.).
[arXiv:0810.3568 [astro-ph]].
[10.1016/j.astropartphys.2008.10.003](https://doi.org/10.1016/j.astropartphys.2008.10.003).
Astropart.Phys. 30 (2009) 293-305.

64) [Upper limits on the VHE gamma-ray emission from the Willman 1 satellite galaxy with the MAGIC Telescope.](#)

By MAGIC Collaboration (E. Aliu et al.).
[arXiv:0810.3561 [astro-ph]].
[10.1088/0004-637X/697/2/1299](https://doi.org/10.1088/0004-637X/697/2/1299).
Astrophys.J. 697 (2009) 1299-1304.

65) [Observation of pulsed gamma-rays above 25 GeV from the Crab pulsar with MAGIC.](#)

By MAGIC Collaboration (E. Aliu et al.).
[arXiv:0809.2998 [astro-ph]].
[10.1126/science.1164718](https://doi.org/10.1126/science.1164718).
Science 322 (2008) 1221-1224.

66) [The TOTEM experiment at the CERN Large Hadron Collider.](#)

By TOTEM Collaboration (G. Anelli et al.).
[10.1088/1748-0221/3/08/S08007](https://doi.org/10.1088/1748-0221/3/08/S08007).
JINST 3 (2008) S08007.

67) [First bounds on the high-energy emission from isolated Wolf-Rayet binary systems.](#)

By MAGIC Collaboration (E. Aliu et al.).
[arXiv:0808.1832 [astro-ph]].
[10.1086/592433](https://doi.org/10.1086/592433).
Astrophys.J. 685 (2008) L71-L74.

68) [Very-High-Energy Gamma Rays from a Distant Quasar: How Transparent Is the Universe?.](#)

By MAGIC Collaboration (E. Aliu et al.).
[arXiv:0807.2822 [astro-ph]].
[10.1126/science.1157087](https://doi.org/10.1126/science.1157087).
Science 320 (2008) 1752, Science 320N5884 (2008) 1752-1754.

69) [Periodic very high energy gamma-ray emission from LS I +61 303 observed with the MAGIC telescope.](#)

By MAGIC Collaboration (J. Albert et al.).
[arXiv:0806.1865 [astro-ph]].
[10.1088/0004-637X/693/1/303](https://doi.org/10.1088/0004-637X/693/1/303).
Astrophys.J. 693 (2009) 303-310.

70) [MAGIC Observations of a 13-Day Flare Complex in M87 in February 2008.](#)

By MAGIC Collaboration (J. Albert et al.).
[arXiv:0806.0988 [astro-ph]].
[10.1086/592348](https://doi.org/10.1086/592348).
Astrophys.J. 685 (2008) L23-L26.

71) [A novel background reduction strategy for high level triggers and processing in gamma-ray Cherenkov detectors.](#)

By MAGIC Collaboration (G. Cabras et al.).
[arXiv:0804.3896 [astro-ph]].
[10.1063/1.2947678](https://doi.org/10.1063/1.2947678).
AIP Conf.Proc. 1018 (2008) 180-183.

72) [Multi-wavelength \(radio, X-ray and gamma-ray\) observations of the gamma-ray binary LS I +61](#)

[303.](#)

By MAGIC Collaboration (J. Albert et al.).

[arXiv:0801.3150 [astro-ph]].

[10.1086/590332.](#)

Astrophys.J. 684 (2008) 1351-1358.

73) [MAGIC observations of the unidentified TeV gamma-ray source TeV J2032+4130.](#)

By MAGIC Collaboration (J. Albert et al.).

[arXiv:0801.2391 [astro-ph]].

[10.1086/529520.](#)

Astrophys.J. 675 (2008) L25-L28.

74) [Frontier detectors for frontier physics. Proceedings, 10th Pisa Meeting on Advanced Detectors, La Biodola, Italy, May 21-27, 2006.](#)

By Franco Cervelli, (ed.), Francesco Forti, (ed.), Riccardo Paoletti, (ed.), Angelo Scribano, (ed.).

75) [The trigger system of the MAGIC telescope.](#)

By R. Paoletti, R. Cecchi, R. Pegna, N. Turini, D. Corti, F. Dazzi, M. Mariotti.

[10.1109/TNS.2007.892649.](#)

IEEE Trans.Nucl.Sci. 54 (2007) 404-409.

76) [A GHz sampling DAQ system for the MAGIC-II telescope.](#)

By Raffaello Pegna, M. Bitossi, R. Cecchi, M. Fagiolini, R. Paoletti, A. Piccioli, N. Turini, M. Barcelo.

[10.1016/j.nima.2006.10.375.](#)

Nucl.Instrum.Meth. A572 (2007) 382-384.

77) [Upper limit for gamma-ray emission above 140-GeV from the dwarf spheroidal galaxy Draco.](#)

By MAGIC Collaboration (J. Albert et al.).

[arXiv:0711.2574 [astro-ph]].

[10.1086/529135.](#)

Astrophys.J. 679 (2008) 428-431.

78) [The MAGIC Project: Contributions to ICRC 2007.](#)

By MAGIC Collaboration (J. Albert et al.).

[arXiv:0709.3763 [astro-ph]].

79) [Implementation of the Random Forest Method for the Imaging Atmospheric Cherenkov Telescope MAGIC.](#)

By J. Albert, E. Aliu, H. Anderhub, P. Antoranz, A. Armada, M. Asensio, C. Baixeras, J.A. Barrio et al..

[arXiv:0709.3719 [astro-ph]].

[10.1016/j.nima.2007.11.068.](#)

Nucl.Instrum.Meth. A588 (2008) 424-432.

80) [The reflecting surface of the MAGIC-II Telescope.](#)

By Denis Bastieri, J. Arnold, C. Baixeras, O. Citterio, F. Dazzi, Barbara De Lotto, M. Doro, M. Ghigo et al..

[arXiv:0709.1372 [astro-ph]].

81) [Probing Quantum Gravity using Photons from a flare of the active galactic nucleus Markarian 501 Observed by the MAGIC telescope.](#)

By MAGIC and Other Contributors Collaborations (J. Albert et al.).

[arXiv:0708.2889 [astro-ph]].

[10.1016/j.physletb.2008.08.053.](#)

Phys.Lett. B668 (2008) 253-257.

82) [Unfolding of differential energy spectra in the MAGIC experiment.](#)

By MAGIC Collaboration (J. Albert et al.).

[arXiv:0707.2453 [astro-ph]].

[10.1016/j.nima.2007.09.048](https://doi.org/10.1016/j.nima.2007.09.048).

Nucl.Instrum.Meth. A583 (2007) 494-506.

83) [Systematic search for VHE gamma-ray emission from X-ray bright high-frequency BL Lac objects.](#)

By MAGIC Collaboration (J. Albert et al.).

[arXiv:0706.4453 [astro-ph]].

[10.1086/587499](https://doi.org/10.1086/587499).

Astrophys.J. 681 (2008) 944-953.

84) [Discovery of Very High Energy gamma-rays from 1ES1011+496 at z=0.212.](#)

By MAGIC Collaboration (J. Albert et al.).

[arXiv:0706.4435 [astro-ph]].

[10.1086/521982](https://doi.org/10.1086/521982).

Astrophys.J. 667 (2007) L21-L23.

85) [Observation of VHE gamma-rays from Cassiopeia A with the MAGIC telescope.](#)

By MAGIC Collaboration (J. Albert et al.).

[arXiv:0706.4065 [astro-ph]].

[10.1051/0004-6361:20078168](https://doi.org/10.1051/0004-6361:20078168).

Astron.Astrophys. 474 (2007) 937-940.

86) [Very High Energy Gamma-ray Radiation from the Stellar-mass Black Hole Cygnus X-1.](#)

By MAGIC Collaboration (J. Albert et al.).

[arXiv:0706.1505 [astro-ph]].

[10.1086/521145](https://doi.org/10.1086/521145).

Astrophys.J. 665 (2007) L51-L54.

87) [VHE Gamma-Ray Observation of the Crab Nebula and Pulsar with MAGIC.](#)

By MAGIC Collaboration (J. Albert et al.).

[arXiv:0705.3244 [astro-ph]].

[10.1086/525270](https://doi.org/10.1086/525270).

Astrophys.J. 674 (2008) 1037-1055.

88) [Discovery of VHE Gamma Radiation from IC443 with the MAGIC Telescope.](#)

By MAGIC Collaboration (J. Albert et al.).

[arXiv:0705.3119 [astro-ph]].

[10.1086/520957](https://doi.org/10.1086/520957).

Astrophys.J. 664 (2007) L87-L90.

89) [Discovery of very high energy gamma-ray emission from the low-frequency peaked BL Lac object BL Lacertae.](#)

By MAGIC Collaboration (J. Albert et al.).

[astro-ph/0703084].

[10.1086/521550](https://doi.org/10.1086/521550).

Astrophys.J. 666 (2007) L17-L20.

90) [Very high energy gamma-ray observations during moonlight and twilight with the MAGIC telescope.](#)

By MAGIC Collaboration (J. Albert et al.).

[astro-ph/0702475].

Submitted to: Astropart.Phys..

91) [Constraints on the steady and pulsed VHE gamma-ray emission from observation of PSR B1951+32/CTB 80 with the MAGIC Telescope.](#)

By MAGIC Collaboration (J. Albert et al.).

[astro-ph/0702077].

[10.1086/521807](#).

Astrophys.J. 669 (2007) 1143-1149.

92) [Variable VHE gamma-ray emission from Markarian 501.](#)

By J. Albert, E. Aliu, H. Anderhub, P. Antoranz, A. Armada, C. Baixeras, J.A. Barrio, H. Bartko et al..
[astro-ph/0702008].

[10.1086/521382](#).

Astrophys.J. 669 (2007) 862-883.

93) [Performance of the Domino Ring Sampler in the MAGIC experiment.](#)

By R. Pegna, R. Cecchi, R. Paoletti, A. Piccioli, N. Turini, S. Ritt.

[10.1016/j.nima.2006.05.205](#).

Nucl.Instrum.Meth. A567 (2006) 218-221.

94) [MAGIC upper limits on the very high energy emission from GRBs.](#)

By MAGIC Collaboration (J. Albert et al.).

[astro-ph/0612548].

[10.1086/520761](#).

Astrophys.J. 667 (2007) 358-366.

95) [FADC signal reconstruction for the MAGIC Telescope.](#)

By MAGIC Collaboration (J. Albert et al.).

[astro-ph/0612385].

[10.1016/j.nima.2008.06.043](#).

Nucl.Instrum.Meth. A594 (2008) 407-419.

96) [Observation of very high energy gamma-rays from the AGN 1ES 2344+514 in a low emission state with the MAGIC telescope.](#)

By MAGIC Collaboration (J. Albert et al.).

[astro-ph/0612383].

[10.1086/518431](#).

Astrophys.J. 662 (2007) 892-899.

97) [First bounds on the very high energy gamma-ray emission from Arp 220.](#)

By MAGIC Collaboration (J. Albert et al.).

[astro-ph/0611786].

[10.1086/511173](#).

Astrophys.J. 658 (2007) 245-248.

98) [Measurement of the \$t \bar{t}\$ Production Cross Section in \$p \bar{p}\$ collisions at \$\sqrt{s} = 1.96\$ -TeV using Lepton + Jets Events with Jet Probability \$b^{\text{-}}\$ tagging.](#)

By CDF Collaboration (A. Abulencia et al.).

[hep-ex/0607035].

[10.1103/PhysRevD.74.072006](#).

Phys.Rev. D74 (2006) 072006.

99) [Observation of \$B^0 \(s\)\$ to \$K^+ K^-\$ and Measurements of Branching Fractions of Charmless Two-body Decays of \$B^0\$ and \$B^0 \(s\)\$ Mesons in \$\bar{p} p\$ Collisions at \$\sqrt{s} = 1.96\$ -TeV.](#)

By CDF Collaboration (A. Abulencia et al.).

[hep-ex/0607021].

[10.1103/PhysRevLett.97.211802](#).

Phys.Rev.Lett. 97 (2006) 211802.

100) [Discovery of very high energy gamma-rays from Markarian 180 triggered by an optical outburst.](#)

By MAGIC Collaboration (J. Albert et al.).

[astro-ph/0606630].

[10.1086/508020](https://arxiv.org/abs/10.1086/508020).

Astrophys.J. 648 (2006) L105-L108.

101) [Search for excited and exotic muons in the \$\mu \gamma\$ decay channel in \$p \bar{p}\$ collisions at \$\sqrt{s} = 1.96\$ -TeV.](#)

By CDF Collaboration (A. Abulencia et al.).

[hep-ex/0606043].

[10.1103/PhysRevLett.97.191802](https://arxiv.org/abs/10.1103/PhysRevLett.97.191802).

Phys.Rev.Lett. 97 (2006) 191802.

102) [Measurement of the \$B^0 \rightarrow B^0_s\$ Oscillation Frequency.](#)

By CDF Collaboration (A. Abulencia et al.).

[hep-ex/0606027].

[10.1103/PhysRevLett.97.062003](https://arxiv.org/abs/10.1103/PhysRevLett.97.062003).

Phys.Rev.Lett. 97 (2006) 062003.

103) [Measurement of the \$t \bar{t}\$ Production Cross Section in \$p \bar{p}\$ Collisions at \$\sqrt{s} = 1.96\$ -TeV.](#)

By CDF Collaboration (A. Abulencia et al.).

[hep-ex/0606017].

[10.1103/PhysRevLett.97.082004](https://arxiv.org/abs/10.1103/PhysRevLett.97.082004).

Phys.Rev.Lett. 97 (2006) 082004.

104) [Detection of VHE radiation from the BL Lac PG 1553+113 with the MAGIC telescope.](#)

By MAGIC Collaboration (J. Albert et al.).

[astro-ph/0606161].

[10.1086/511384](https://arxiv.org/abs/10.1086/511384).

Astrophys.J. 654 (2007) L119-L122.

105) [Search for a neutral Higgs boson decaying to a \$W\$ boson pair in \$p \bar{p}\$ collisions at \$\sqrt{s} = 1.96\$ -TeV.](#)

By CDF Collaboration (A. Abulencia et al.).

[hep-ex/0605124].

[10.1103/PhysRevLett.97.081802](https://arxiv.org/abs/10.1103/PhysRevLett.97.081802).

Phys.Rev.Lett. 97 (2006) 081802.

106) [Top quark mass measurement from dilepton events at CDF II with the matrix-element method.](#)

By CDF Collaboration (A. Abulencia et al.).

[hep-ex/0605118].

[10.1103/PhysRevD.74.032009](https://arxiv.org/abs/10.1103/PhysRevD.74.032009).

Phys.Rev. D74 (2006) 032009.

107) [Measurement of the \$b\$ jet cross-section in events with a \$Z\$ boson in \$p \bar{p}\$ collisions at \$\sqrt{s} = 1.96\$ -TeV.](#)

By CDF Collaboration (A. Abulencia et al.).

[hep-ex/0605099].

[10.1103/PhysRevD.74.032008](https://arxiv.org/abs/10.1103/PhysRevD.74.032008).

Phys.Rev. D74 (2006) 032008.

108) [Search for new physics in lepton + photon + \$X\$ events with 305 \$\text{pb}^{-1}\$ of \$p \bar{p}\$ collisions at \$\sqrt{s} = 1.96\$ -TeV.](#)

By CDF Collaboration (A. Abulencia et al.).

[hep-ex/0605097].

[10.1103/PhysRevLett.97.031801](https://arxiv.org/abs/10.1103/PhysRevLett.97.031801).

Phys.Rev.Lett. 97 (2006) 031801.

- 109) [Variable Very High Energy Gamma-ray Emission from the Microquasar LS I +61 303.](#)
By MAGIC Collaboration (J. Albert et al.).
[astro-ph/0605549].
[10.1126/SCIENCE.1128177.](#)
Science 312 (2006) 1771-1773.
- 110) [A PMT-block test bench.](#)
By P. Adragna, A. Antonaki, I. Boudagov, V. Cavasinni, D. Costanzo, T. Del Prete, A. Dotti, D. Fassouliotis et al..
[physics/0605074].
[10.1016/j.nima.2006.03.045.](#)
Nucl.Instrum.Meth. A564 (2006) 597-607.
- 111) [Measurement of the ratio of branching fractions \$B\(D0 \to K^{+} \pi^{-}\) / B\(D0 \to K^{-} \pi^{+}\)\$ using the CDF II Detector.](#)
By CDF Collaboration (A. Abulencia et al.).
[hep-ex/0605027].
[10.1103/PhysRevD.74.031109.](#)
Phys.Rev. D74 (2006) 031109.
- 112) [Search for Large Extra Dimensions in the Production of Jets and Missing Transverse Energy in p anti-p Collisions at \$s^{1/2} = 1.96\$ TeV.](#)
By CDF Collaboration (A. Abulencia et al.).
[hep-ex/0605101].
[10.1103/PhysRevLett.97.171802.](#)
Phys.Rev.Lett. 97 (2006) 171802.
- 113) [Observation of VHE Gamma Radiation from HESS J1834-087/W41 with the MAGIC Telescope.](#)
By MAGIC Collaboration (J. Albert et al.).
[astro-ph/0604197].
[10.1086/504917.](#)
Astrophys.J. 643 (2006) L53-L56.
- 114) [Measurement of the t anti-t production cross section in p anti-p collisions at \$s^{1/2} = 1.96\$ -TeV using missing \$E\(T\) +\$ jets events with secondary vertex b-tagging.](#)
By CDF Collaboration (A. Abulencia et al.).
[hep-ex/0603043].
[10.1103/PhysRevLett.96.202002.](#)
Phys.Rev.Lett. 96 (2006) 202002.
- 115) [Discovery of VHE gamma-ray emission from 1ES1218+30.4.](#)
By MAGIC Collaboration (J. Albert et al.).
[astro-ph/0603529].
[10.1086/504845.](#)
Astrophys.J. 642 (2006) L119-L122.
- 116) [Observations of mkn 421 with the magic telescope.](#)
By MAGIC Collaboration (J. Albert et al.).
[astro-ph/0603478].
[10.1086/518221.](#)
Astrophys.J. 663 (2007) 125-138.
- 117) [Measurement of the B\(c\)+ meson lifetime using \$B\(c\)^+ \to J/\psi e^+ \nu\(e\)\$.](#)
By CDF Collaboration (A. Abulencia et al.).
[hep-ex/0603027].

[10.1103/PhysRevLett.97.012002.](#)
Phys.Rev.Lett. 97 (2006) 012002.

118) [Search for high-mass resonances decaying to \$e\mu\$ in \$p\bar{p}\$ collisions at \$\sqrt{s} = 1.96\$ -TeV..](#)

By CDF Collaboration (A. Abulencia et al.).
[hep-ex/0603006].

[10.1103/PhysRevLett.96.211802.](#)
Phys.Rev.Lett. 96 (2006) 211802.

119) [Search for \$Z^{\prime} \rightarrow e^+e^-\$ using dielectron mass and angular distribution..](#)

By CDF Collaboration (A. Abulencia et al.).
[hep-ex/0602045].

[10.1103/PhysRevLett.96.211801.](#)
Phys.Rev.Lett. 96 (2006) 211801.

120) [TOTEM physics.](#)

By TOTEM Collaboration (G. Anelli et al.).
[hep-ex/0602025].

121) [Flux upper limit of gamma-ray emission by grb050713a from magic telescope observations.](#)

By J. Albert, E. Aliu, H. Anderhub, P. Antoranz, A. Armada, M. Asensio, C. Baixeras, J.A. Barrio et al..
[astro-ph/0602231].

[10.1086/503767.](#)

Astrophys.J. 641 (2006) L9-L12.

122) [Diffraction and total cross-section at the Tevatron and the LHC.](#)

By TOTEM Collaboration (M. Deile et al.).
[hep-ex/0602021].

123) [Measurement of the top quark mass using template methods on dilepton events in proton antiproton collisions at \$\sqrt{s} = 1.96\$ -TeV..](#)

By CDF Collaboration (A. Abulencia et al.).
[hep-ex/0602008].

[10.1103/PhysRevD.73.112006.](#)

Phys.Rev. D73 (2006) 112006.

124) [Observation of \$B_s^0 \rightarrow \psi\(2S\) \phi\$ and measurement of ratio of branching fractions \$\frac{B\(B_s^0 \rightarrow \psi\(2S\) \phi\)}{B\(B_s^0 \rightarrow J/\psi \phi\)}\$.](#)

By CDF Collaboration (A. Abulencia et al.).
[hep-ex/0602005].

[10.1103/PhysRevLett.96.231801.](#)

Phys.Rev.Lett. 96 (2006) 231801.

125) [Measurement of \$\frac{\sigma\(\Lambda_b^0\) \times \text{BR}\(\Lambda_b^0 \rightarrow \Lambda_c^+ \pi^-\)}{\sigma\(\bar{B}^0\) \times \text{BR}\(\bar{B}^0 \rightarrow D^+ \pi^-\)}\$ in \$p\bar{p}\$ collisions at \$\sqrt{s} = 1.96\$ -TeV..](#)

By CDF Collaboration (A. Abulencia et al.).
[hep-ex/0601003].

[10.1103/PhysRevLett.98.122002.](#)

Phys.Rev.Lett. 98 (2007) 122002.

126) [A third level trigger programmable on FPGA for the gamma / hadron separation in a Cherenkov telescope using Pseudo-Zernike moments and the SVM classifier.](#)

By M. Frailis, O. Mansutti, P. Boinee, G. Cabras, A. De Angelis, B. De Lotto, A. Forti, M. Dell'Orso et al..

- 127) [Measurement of the dipion mass spectrum in \$X\(3872\) \rightarrow J/\psi \pi^+ \pi^-\$ decays.](#)
By CDF Collaboration (A. Abulencia et al.).
[hep-ex/0512074].
[10.1103/PhysRevLett.96.102002](#).
Phys.Rev.Lett. 96 (2006) 102002.
- 128) [A search for scalar bottom quarks from gluino decays in \$\bar{p}p\$ collisions at \$\sqrt{s} = 1.96\$ -TeV.](#)
By CDF Collaboration (A. Abulencia et al.).
[hep-ex/0512072].
[10.1103/PhysRevLett.96.171802](#).
Phys.Rev.Lett. 96 (2006) 171802.
- 129) [Top quark mass measurement from dilepton events at CDF II.](#)
By CDF Collaboration (A. Abulencia et al.).
[hep-ex/0512070].
[10.1103/PhysRevLett.96.152002](#).
Phys.Rev.Lett. 96 (2006) 152002.
- 130) [Measurement of mass and width of the excited charmed meson states \$D_0\(1\)\$ and \$D^*\(2\)\$ at CDF.](#)
By CDF Collaboration (A. Abulencia et al.).
[hep-ex/0512069].
[10.1103/PhysRevD.73.051104](#).
Phys.Rev. D73 (2006) 051104.
- 131) [Search for anomalous semileptonic decay of heavy flavor hadrons produced in association with a \$W\$ boson at CDF II.](#)
By CDF Collaboration (A. Abulencia et al.).
[hep-ex/0512065].
[10.1103/PhysRevD.73.051101](#).
Phys.Rev. D73 (2006) 051101.
- 132) [Measurement of the inclusive jet cross section using the \$k\(t\)\$ algorithm in \$p\$ anti- \$p\$ collisions at \$\sqrt{s} = 1.96\$ -TeV.](#)
By CDF Collaboration (A. Abulencia et al.).
[hep-ex/0512062].
[10.1103/PhysRevLett.96.122001](#).
Phys.Rev.Lett. 96 (2006) 122001.
- 133) [Search for \$H\$ to \$b\$ anti- \$b\$ produced in association with \$W\$ bosons in \$p\bar{p}\$ collisions at \$\sqrt{s} = 1.96\$ -TeV.](#)
By CDF Collaboration (A. Abulencia et al.).
[hep-ex/0512051].
[10.1103/PhysRevLett.96.081803](#).
Phys.Rev.Lett. 96 (2006) 081803.
- 134) [Observation of gamma-rays from the galactic center with the magic telescope.](#)
By MAGIC Collaboration (J. Albert et al.).
[astro-ph/0512469].
[10.1086/501164](#).
Astrophys.J. 638 (2006) L101-L104.
- 135) [Search for second-generation scalar leptoquarks in \$p\bar{p}\$ collisions at \$\sqrt{s} = 1.96\$ -TeV.](#)
By CDF Collaboration (A. Abulencia et al.).
[hep-ex/0512055].

[10.1103/PhysRevD.73.051102.](#)

Phys.Rev. D73 (2006) 051102.

136) [Magic observations of very high energy gamma-rays from HESS J1813-178.](#)

By MAGIC Collaboration (J. Albert et al.).

[astro-ph/0512283].

[10.1086/500364.](#)

Astrophys.J. 637 (2006) L41-L44.

137) [Measurement of the inclusive jet cross section in \$p\bar{p}\$ interactions at \$\sqrt{s} = 1.96\$ -TeV using a cone-based jet algorithm..](#)

By CDF Collaboration (A. Abulencia et al.).

[hep-ex/0512020].

[10.1103/PhysRevD.74.071103.](#)

Phys.Rev. D74 (2006) 071103.

138) [Measurement of the top quark mass with the dynamical likelihood method using lepton plus jets events with b-tags in \$p\bar{p}\$ collisions at \$\sqrt{s} = 1.96\$ -TeV..](#)

By CDF Collaboration (A. Abulencia et al.).

[hep-ex/0512009].

[10.1103/PhysRevD.73.092002.](#)

Phys.Rev. D73 (2006) 092002.

139) [Measurement of the helicity of \$W\$ bosons in top-quark decays.](#)

By CDF Collaboration (A. Abulencia et al.).

[hep-ex/0511023].

[10.1103/PhysRevD.73.111103.](#)

Phys.Rev. D73 (2006) 111103.

140) [Search for charged Higgs bosons from top quark decays in \$p\bar{p}\$ collisions at \$\sqrt{s} = 1.96\$ -TeV..](#)

By CDF Collaboration (A. Abulencia et al.).

[hep-ex/0510065].

[10.1103/PhysRevLett.96.042003.](#)

Phys.Rev.Lett. 96 (2006) 042003.

141) [A search for \$t \rightarrow \tau \nu q\$ in \$t\bar{t}\$ production..](#)

By CDF Collaboration (A. Abulencia et al.).

[hep-ex/0510063].

[10.1016/j.physletb.2006.06.030.](#)

Phys.Lett. B639 (2006) 172-178.

142) [Technical performance of the MAGIC telescope.](#)

By MAGIC Collaboration (J. Cortina et al.).

143) [The data acquisition of the MAGIC II telescope.](#)

By MAGIC Collaboration (Jose Antonio Coarasa Perez et al.).

144) [MAGIC phase II.](#)

By MAGIC Collaboration (C. Baixeras et al.).

145) [Precision top quark mass measurement in the lepton + jets topology in \$p\$ anti- \$p\$ collisions at \$\sqrt{s} = 1.96\$ -TeV..](#)

By CDF Collaboration (A. Abulencia et al.).

[hep-ex/0510049].

[10.1103/PhysRevLett.96.022004.](#)

Phys.Rev.Lett. 96 (2006) 022004.

146) [Top quark mass measurement using the template method in the lepton + jets channel at CDF II.](#)

By CDF Collaboration (A. Abulencia et al.).

[hep-ex/0510048].

[10.1103/PhysRevD.73.032003.](#)

Phys.Rev. D73 (2006) 032003.

147) [Direct search for Dirac magnetic monopoles in \$p\bar{p}\$ collisions at \$\sqrt{s} = 1.96\$ TeV.](#)

By CDF Collaboration (A. Abulencia et al.).

[hep-ex/0509015].

[10.1103/PhysRevLett.96.201801.](#)

Phys.Rev.Lett. 96 (2006) 201801.

148) [Search for neutral MSSM Higgs bosons decaying to tau pairs in \$p\bar{p}\$ collisions at \$\sqrt{s} = 1.96\$ TeV.](#)

By CDF Collaboration (A. Abulencia et al.).

[hep-ex/0508051].

[10.1103/PhysRevLett.96.011802.](#)

Phys.Rev.Lett. 96 (2006) 011802.

149) [Observation of VHE gamma-ray emission from the active galactic nucleus 1ES1959+650 using the Magic Telescope.](#)

By MAGIC Collaboration (J. Albert et al.).

[astro-ph/0508543].

[10.1086/499421.](#)

Astrophys.J. 639 (2006) 761-765.

150) [Search for \$B_s \rightarrow \mu^+ \mu^-\$ and \$B_d \rightarrow \mu^+ \mu^-\$ decays in \$p\bar{p}\$ collisions with CDF II.](#)

By CDF Collaboration (A. Abulencia et al.).

[hep-ex/0508036].

[10.1103/PhysRevLett.95.221805.](#)

Phys.Rev.Lett. 95 (2005) 221805, Erratum-ibid. 95 (2005) 249905.

151) [The MAGIC project: Contributions to ICRC 2005, Pune, India. Part 1.](#)

By MAGIC Collaboration (J. Albert i Fort et al.).

[astro-ph/0508244].

152) [Measurements of inclusive W and Z cross sections in p anti-p collisions at \$s^{*1/2} = 1.96\$ -TeV.](#)

By CDF Collaboration (A. Abulencia et al.).

[hep-ex/0508029].

[10.1088/0954-3899/34/12/001.](#)

J.Phys. G34 (2007) 2457-2544.

153) [Measurement of \$B\$ hadron masses in exclusive \$J/\psi\$ decays with the CDF detector.](#)

By CDF Collaboration (D. Acosta et al.).

[hep-ex/0508022].

[10.1103/PhysRevLett.96.202001.](#)

Phys.Rev.Lett. 96 (2006) 202001.

154) [Measurement of the ratios of branching fractions \$B\(B^0 \rightarrow D^+ \pi^-\)/B\(B^0 \rightarrow D^+ \pi^0\)\$ and \$B\(B^+ \rightarrow \overline{D}^0 \pi^+\)/B\(B^+ \rightarrow D^+ \pi^0\)\$.](#)

By CDF Collaboration (A. Abulencia et al.).

[hep-ex/0508014].

[10.1103/PhysRevLett.96.191801.](#)

Phys.Rev.Lett. 96 (2006) 191801.

155) [Search for new high mass particles decaying to lepton pairs in \$p\bar{p}\$ collisions at \$\sqrt{s} = 1.96\$ TeV.](#)

By CDF Collaboration (A. Abulencia et al.).
[hep-ex/0507104].

[10.1103/PhysRevLett.95.252001.](#)

Phys.Rev.Lett. 95 (2005) 252001.

156) [Physics and astrophysics with a ground-based gamma-ray telescope of low energy threshold.](#)

By J. Albert i Fort, A. Armada, C. Baixeras, H. Bartko, D. Bastieri, W. Bednarek, C. Bigongiari, A. Biland et al..

[10.1016/j.astropartphys.2005.03.005.](#)

Astropart.Phys. 23 (2005) 493-509.

157) [Search for \$\Lambda_b \rightarrow p \pi\$ and \$\Lambda_b \rightarrow p K\$ decays in \$p\bar{p}\$ collisions at \$\sqrt{s} = 1.96\$ TeV.](#)

By CDF Collaboration (D. Acosta et al.).
[hep-ex/0507067].

[10.1103/PhysRevD.72.051104.](#)

Phys.Rev. D72 (2005) 051104.

158) [Search for W and Z bosons in the reaction \$p\bar{p} \rightarrow 2 \text{ jets} + \gamma\$ at \$\sqrt{s} = 1.8\$ TeV.](#)

By CDF Collaboration (D. Acosta et al.).
[hep-ex/0507051].

[10.1103/PhysRevD.73.012001.](#)

Phys.Rev. D73 (2006) 012001.

159) [Search for first-generation scalar leptoquarks in \$p\bar{p}\$ collisions at \$\sqrt{s} = 1.96\$ TeV.](#)

By CDF Collaboration (D. Acosta et al.).
[hep-ex/0506074].

[10.1103/PhysRevD.72.051107.](#)

Phys.Rev. D72 (2005) 051107.

160) [A search for supersymmetric Higgs bosons in the di-tau decay mode in \$p\bar{p}\$ collisions at \$\sqrt{s} = 1.8\$ TeV.](#)

By CDF Collaboration (D. Acosta et al.).
[hep-ex/0506042].

[10.1103/PhysRevD.72.072004.](#)

Phys.Rev. D72 (2005) 072004.

161) [Search for new physics using high mass tau pairs from 1.96 TeV \$p\bar{p}\$ collisions.](#)

By CDF Collaboration (D. Acosta et al.).
[hep-ex/0506034].

[10.1103/PhysRevLett.95.131801.](#)

Phys.Rev.Lett. 95 (2005) 131801.

162) [Measurement of the \$t\bar{t}\$ production cross section in \$p\bar{p}\$ collisions at \$\sqrt{s} = 1.96\$ TeV using lepton plus jets events with semileptonic B decays to muons.](#)

By CDF Collaboration (D. Acosta et al.).
[hep-ex/0506001].

[10.1103/PhysRevD.72.032002.](#)

Phys.Rev. D72 (2005) 032002.

163) [Measurement of \$B\(t \rightarrow W b\)/B\(t \rightarrow W q\)\$ at the Collider Detector at Fermilab.](#)

By CDF Collaboration (D. Acosta et al.).
[hep-ex/0505091].
[10.1103/PhysRevLett.95.102002](https://arxiv.org/abs/10.1103/PhysRevLett.95.102002).
Phys.Rev.Lett. 95 (2005) 102002.

164) [Evidence for the exclusive decay \$B_c^{\pm} \rightarrow J/\psi \pi^{\pm}\$ and measurement of the mass of the \$B_c\$ meson.](#)

By CDF Collaboration (A. Abulencia et al.).
[hep-ex/0505076].
[10.1103/PhysRevLett.96.082002](https://arxiv.org/abs/10.1103/PhysRevLett.96.082002).
Phys.Rev.Lett. 96 (2006) 082002.

165) [Study of jet shapes in inclusive jet production in \$p\bar{p}\$ collisions at \$\sqrt{s}=1.96\$ TeV.](#)

By CDF Collaboration (D. Acosta et al.).
[hep-ex/0505013].
[10.1103/PhysRevD.71.112002](https://arxiv.org/abs/10.1103/PhysRevD.71.112002).
Phys.Rev. D71 (2005) 112002.

166) [Measurement of the cross section for \$t\bar{t}\$ production in \$p\bar{p}\$ collisions using the kinematics of lepton + jets events.](#)

By CDF Collaboration (D. Acosta et al.).
[hep-ex/0504053].
[10.1103/PhysRevD.72.052003](https://arxiv.org/abs/10.1103/PhysRevD.72.052003).
Phys.Rev. D72 (2005) 052003.

167) [\$K^0_S\$ and \$\Lambda^0\$ production studies in \$p\bar{p}\$ collisions at \$\sqrt{s}=1800\$ -GeV and 630-GeV.](#)

By CDF Collaboration (D. Acosta et al.).
[hep-ex/0504048].
[10.1103/PhysRevD.72.052001](https://arxiv.org/abs/10.1103/PhysRevD.72.052001).
Phys.Rev. D72 (2005) 052001.

168) [Measurement of the azimuthal angle distribution of leptons from \$W\$ boson decays as a function of the \$W\$ transverse momentum in \$p\bar{p}\$ collisions at \$\sqrt{s} = 1.8\$ TeV.](#)

By CDF Collaboration (D. Acosta et al.).
[hep-ex/0504020].
[10.1103/PhysRevD.73.052002](https://arxiv.org/abs/10.1103/PhysRevD.73.052002).
Phys.Rev. D73 (2006) 052002.

169) [Search for Higgs bosons decaying into \$b\bar{b}\$ and produced in association with a vector boson in \$p\bar{p}\$ collisions at \$\sqrt{s} = 1.8\$ TeV.](#)

By CDF Collaboration (D. Acosta et al.).
[hep-ex/0503039].
[10.1103/PhysRevLett.95.051801](https://arxiv.org/abs/10.1103/PhysRevLett.95.051801).
Phys.Rev.Lett. 95 (2005) 051801.

170) [Search for long-lived doubly-charged Higgs bosons in \$p\bar{p}\$ collisions at \$\sqrt{s} = 1.96\$ TeV.](#)

By CDF Collaboration (D. Acosta et al.).
[hep-ex/0503004].
[10.1103/PhysRevLett.95.071801](https://arxiv.org/abs/10.1103/PhysRevLett.95.071801).
Phys.Rev.Lett. 95 (2005) 071801.

171) [First evidence for \$B_s^0 \rightarrow \psi \psi\$ decay and measurements of branching ratio and \$A_{CP}\$ for \$B^+ \rightarrow \psi K^+\$.](#)

By CDF Collaboration (D. Acosta et al.).

[hep-ex/0502044].

[10.1103/PhysRevLett.95.031801](#).

Phys.Rev.Lett. 95 (2005) 031801.

172) [Measurement of the moments of the hadronic invariant mass distribution in semileptonic \$B_s\$ decays.](#)

By CDF Collaboration (D. Acosta et al.).

[hep-ex/0502003].

[10.1103/PhysRevD.71.051103](#).

Phys.Rev. D71 (2005) 051103.

173) [Measurement of the \$W^+ W^-\$ production cross section in \$p\bar{p}\$ collisions at \$\sqrt{s} = 1.96\$ TeV using dilepton events.](#)

By CDF Collaboration (D. Acosta et al.).

[hep-ex/0501050].

[10.1103/PhysRevLett.94.211801](#).

Phys.Rev.Lett. 94 (2005) 211801.

174) [Measurement of the forward-backward charge asymmetry from \$W \rightarrow e \nu\$ production in \$p\bar{p}\$ collisions at \$\sqrt{s} = 1.96\$ TeV.](#)

By CDF Collaboration (D. Acosta et al.).

[hep-ex/0501023].

[10.1103/PhysRevD.71.051104](#).

Phys.Rev. D71 (2005) 051104.

175) [Search for \$Z Z\$ and \$Z W\$ production in \$p\bar{p}\$ collisions at \$\sqrt{s} = 1.96\$ TeV.](#)

By CDF Collaboration (D. Acosta et al.).

[hep-ex/0501021].

[10.1103/PhysRevD.71.091105](#).

Phys.Rev. D71 (2005) 091105.

176) [First results from MAGIC.](#)

By MAGIC Collaboration (R. Paoletti for the collaboration).

177) [Performance of a 3D imaging electromagnetic calorimeter for the AMS02 space experiment.](#)

By C. Adloff, G. Coignet, L. Girard, C. Goy, R. Kossakowski, S. Rosier-Lees, J. Pochon, J.P. Vialle et al..

178) [Measurement of the \$J/\psi\$ meson and \$b\$ -hadron production cross sections in \$p\bar{p}\$ collisions at \$\sqrt{s} = 1960\$ GeV.](#)

By CDF Collaboration (D. Acosta et al.).

[hep-ex/0412071].

[10.1103/PhysRevD.71.032001](#).

Phys.Rev. D71 (2005) 032001.

179) [Measurement of the lifetime difference between \$B\(s\)\$ mass eigenstates.](#)

By CDF Collaboration (D. Acosta et al.).

[hep-ex/0412057].

[10.1103/PhysRevLett.94.101803](#).

Phys.Rev.Lett. 94 (2005) 101803.

180) [Measurement of the cross section for prompt diphoton production in \$p\bar{p}\$ collisions at \$\sqrt{s} = 1.96\$ TeV.](#)

By CDF Collaboration (D. Acosta et al.).

[hep-ex/0412050].

[10.1103/PhysRevLett.95.022003](#).

Phys.Rev.Lett. 95 (2005) 022003.

- 181) [Search for anomalous kinematics in \$t\bar{t}\$ dilepton events at {CDF} II.](#)
By CDF Collaboration (D. Acosta et al.).
[hep-ex/0412042].
[10.1103/PhysRevLett.95.022001](#).
Phys.Rev.Lett. 95 (2005) 022001.
- 182) [Measurements of \$b\bar{b}\$ azimuthal production correlations in \$p\bar{p}\$ collisions at \$\sqrt{s} = 1.8\$ TeV.](#)
By CDF Collaboration (D. Acosta et al.).
[hep-ex/0412006].
[10.1103/PhysRevD.71.092001](#).
Phys.Rev. D71 (2005) 092001.
- 183) [Measurement of the \$W\$ boson polarization in top decay at {CDF} at \$\sqrt{s} = 1.8\$ TeV.](#)
By CDF Collaboration (D. Acosta et al.).
[hep-ex/0411070].
[10.1103/PhysRevD.71.031101](#), [10.1103/PhysRevD.71.059901](#).
Phys.Rev. D71 (2005) 031101, Erratum-ibid. D71 (2005) 059901.
- 184) [Measurement of the forward-backward charge asymmetry of electron positron pairs in \$p\bar{p}\$ collisions at \$\sqrt{s} = 1.96\$ TeV.](#)
By CDF Collaboration (D. Acosta et al.).
[hep-ex/0411059].
[10.1103/PhysRevD.71.052002](#).
Phys.Rev. D71 (2005) 052002.
- 185) [Measurement of charged particle multiplicities in gluon and quark jets in \$p\bar{p}\$ collisions at \$\sqrt{s} = 1.8\$ TeV.](#)
By CDF Collaboration (D. Acosta et al.).
[10.1103/PhysRevLett.94.171802](#).
Phys.Rev.Lett. 94 (2005) 171802.
- 186) [Search for scalar leptoquark pairs decaying to \$\nu\bar{\nu} q\bar{q}\$ in \$p\bar{p}\$ collisions at \$\sqrt{s} = 1.96\$ TeV.](#)
By CDF Collaboration (D. Acosta et al.).
[hep-ex/0410076].
[10.1103/PhysRevD.71.119901](#), [10.1103/PhysRevD.71.112001](#).
Phys.Rev. D71 (2005) 112001, Erratum-ibid. D71 (2005) 119901.
- 187) [Search for electroweak single top quark production in \$p\bar{p}\$ collisions at \$\sqrt{s} = 1.96\$ TeV.](#)
By CDF Collaboration (D. Acosta et al.).
[hep-ex/0410058].
[10.1103/PhysRevD.71.012005](#).
Phys.Rev. D71 (2005) 012005.
- 188) [Search for anomalous production of diphoton events with missing transverse energy at CDF and limits on gauge-mediated supersymmetry-breaking models.](#)
By CDF Collaboration (D. Acosta et al.).
[hep-ex/0410053].
[10.1103/PhysRevD.71.031104](#).
Phys.Rev. D71 (2005) 031104.
- 189) [Measurement of the \$t\bar{t}\$ production cross section in \$p\bar{p}\$ collisions at \$\sqrt{s} = 1.96\$ TeV using lepton + jets events with secondary vertex \$b\$ -tagging.](#)

By CDF Collaboration (D. Acosta et al.).
[hep-ex/0410041].
[10.1103/PhysRevD.71.052003.](#)
Phys.Rev. D71 (2005) 052003.

190) [Comparison of three-jet events in \$p\bar{p}\$ collisions at \$\sqrt{s} = 1.8\$ TeV to predictions from a next-to-leading order QCD calculation.](#)
By CDF Collaboration (D. Acosta et al.).
[hep-ex/0410018].
[10.1103/PhysRevD.71.032002.](#)
Phys.Rev. D71 (2005) 032002.

191) [Measurement of partial widths and search for direct CP violation in \$D^0\$ meson decays to \$K^+K^-\$ and \$\pi^+\pi^-\$.](#)
By CDF Collaboration (D. Acosta et al.).
[hep-ex/0504006].
[10.1103/PhysRevLett.94.122001.](#)
Phys.Rev.Lett. 94 (2005) 122001.

192) [Search for excited and exotic electrons in the \$e\gamma\$ decay channel in \$p\bar{p}\$ collisions at \$\sqrt{s} = 1.96\$ TeV.](#)
By CDF Collaboration (D. Acosta et al.).
[hep-ex/0410013].
[10.1103/PhysRevLett.94.101802.](#)
Phys.Rev.Lett. 94 (2005) 101802.

193) [Measurement of \$W\gamma\$ and \$Z\gamma\$ production in \$p\bar{p}\$ collisions at \$\sqrt{s} = 1.96\$ TeV.](#)
By CDF Collaboration (D. Acosta et al.).
[hep-ex/0410008].
[10.1103/PhysRevLett.94.041803.](#)
Phys.Rev.Lett. 94 (2005) 041803.

194) [Measurement of the \$t\bar{t}\$ production cross section in \$p\bar{p}\$ collisions at \$\sqrt{s} = 1.96\$ TeV using kinematic fitting of \$b\$ -tagged lepton + jet events.](#)
By CDF Collaboration (D. Acosta et al.).
[hep-ex/0409029].
[10.1103/PhysRevD.71.072005.](#)
Phys.Rev. D71 (2005) 072005.

195) [First measurements of inclusive \$W\$ and \$Z\$ cross sections from Run II of the Tevatron collider.](#)
By CDF Collaboration (D. Acosta et al.).
[hep-ex/0406078].
[10.1103/PhysRevLett.94.091803.](#)
Phys.Rev.Lett. 94 (2005) 091803.

196) [Search for doubly-charged Higgs bosons decaying to dileptons in \$p\bar{p}\$ collisions at \$\sqrt{s} = 1.96\$ TeV.](#)
By CDF Collaboration (D. Acosta et al.).
[hep-ex/0406073].
[10.1103/PhysRevLett.93.221802.](#)
Phys.Rev.Lett. 93 (2004) 221802.

197) [Inclusive search for anomalous production of high \$p_T\$ like-sign lepton pairs in \$p\bar{p}\$ collisions at \$\sqrt{s} = 1.8\$ TeV.](#)
By CDF Collaboration (D. Acosta et al.).

[hep-ex/0405063].

[10.1103/PhysRevLett.93.061802](https://doi.org/10.1103/PhysRevLett.93.061802).

Phys.Rev.Lett. 93 (2004) 061802.

198) [Measurement of the \$t\bar{t}\$ production cross section in \$p\bar{p}\$ collisions at \$\sqrt{s} = 1.96\$ TeV using dilepton events.](#)

By CDF Collaboration (D. Acosta et al.).

[hep-ex/0404036].

[10.1103/PhysRevLett.93.142001](https://doi.org/10.1103/PhysRevLett.93.142001).

Phys.Rev.Lett. 93 (2004) 142001.

199) [Direct photon cross section with conversions at CDF.](#)

By CDF Collaboration (D. Acosta et al.).

[hep-ex/0404022].

[10.1103/PhysRevD.70.074008](https://doi.org/10.1103/PhysRevD.70.074008).

Phys.Rev. D70 (2004) 074008.

200) [The trigger system of the MAGIC telescope: On-line selection strategies for Cherenkov telescopes.](#)

By MAGIC Collaboration (M. Meucci et al.).

[10.1016/j.nima.2003.11.083](https://doi.org/10.1016/j.nima.2003.11.083).

Nucl.Instrum.Meth. A518 (2004) 554-556.

201) [Commissioning and first tests of the MAGIC telescope.](#)

By C. Baixeras, D. Bastieri, C. Bigongiari, O. Blanch, G. Blanchot, R. Bock, T. Bretz, A. Chilingarian et al..

[10.1016/j.nima.2003.10.057](https://doi.org/10.1016/j.nima.2003.10.057).

Nucl.Instrum.Meth. A518 (2004) 188-192.

202) [The underlying event in hard interactions at the Tevatron \$p\bar{p}\$ collider.](#)

By CDF Collaboration (D. Acosta et al.).

[hep-ex/0404004].

[10.1103/PhysRevD.70.072002](https://doi.org/10.1103/PhysRevD.70.072002).

Phys.Rev. D70 (2004) 072002.

203) [Optimized search for single top quark production at the Fermilab tevatron.](#)

By CDF Collaboration (D. Acosta et al.).

[10.1103/PhysRevD.69.052003](https://doi.org/10.1103/PhysRevD.69.052003).

Phys.Rev. D69 (2004) 052003.

204) [Search for \$B_s^0 \rightarrow \mu^+ \mu^-\$ and \$B_d^0 \rightarrow \mu^+ \mu^-\$ decays in \$p\bar{p}\$ collisions at \$\sqrt{s} = 1.96\$ TeV.](#)

By CDF Collaboration (D. Acosta et al.).

[hep-ex/0403032].

[10.1103/PhysRevLett.93.032001](https://doi.org/10.1103/PhysRevLett.93.032001).

Phys.Rev.Lett. 93 (2004) 032001.

205) [Design studies for ECO, the European gamma-ray observatory.](#)

By Carmen Baixeras, Denis Bastieri, Wlodek Bednarek, Ciro Bigongiari, Adrian Biland, Oscar Blanch, Rudolf K. Bock, Thomas Bretz et al..

[astro-ph/0403180].

206) [Performances of the AMS-02 electromagnetic calorimeter.](#)

By L. Girard, C. Adloff, G. Coignet, C. Goy, R. Kossakowski, S. Rosier-Lees, J. Pochon, J.P. Vialle et al..

207) [Technical innovations for the MAGIC project.](#)

By MAGIC Collaboration (R. Mirzorian et al.).

208) [The trigger system of the MAGIC telescope: On-line selection strategies for Cherenkov telescopes.](#)
By MAGIC Collaboration (D. Corti et al.).

209) [Extending the Cherenkov technique down to an energy threshold of a few GeV: The ultimate instrument for ground-based gamma-ray astronomy.](#)
By M. Merck, C. Baixeras, A. Biland, J. Cortina, D. Ferenc, M.V. Fonseca, E. Lorenz, M. Mariotti et al..

210) [Observation of the narrow state \$X\(3872\) \rightarrow J/\psi \pi^+ \pi^-\$ in \$\bar{p}p\$ collisions at \$\sqrt{s} = 1.96\$ TeV.](#)
By CDF Collaboration (D. Acosta et al.).
[hep-ex/0312021].
[10.1103/PhysRevLett.93.072001.](#)
Phys.Rev.Lett. 93 (2004) 072001.

211) [Heavy flavor properties of jets produced in \$\bar{p}p\$ interactions at \$\sqrt{s} = 1.8\$ -TeV.](#)
By CDF Collaboration (D. Acosta et al.).
[hep-ex/0311051].
[10.1103/PhysRevD.69.072004.](#)
Phys.Rev. D69 (2004) 072004.

212) [Measurement of the polar-angle distribution of leptons from \$W\$ boson decay as a function of the \$W\$ transverse momentum in \$\bar{p}p\$ collisions at \$\sqrt{s} = 1.8\$ TeV.](#)
By CDF Collaboration (D. Acosta et al.).
[hep-ex/0311050].
[10.1103/PhysRevD.70.032004.](#)
Phys.Rev. D70 (2004) 032004.

213) [Combination of {CDF} and D\O results on \$W\$ boson mass and width.](#)
By CDF and D0 Collaborations (V.M. Abazov et al.).
[hep-ex/0311039].
[10.1103/PhysRevD.70.092008.](#)
Phys.Rev. D70 (2004) 092008.

214) [Inclusive double pomeron exchange at the Fermilab Tevatron \$\bar{p}p\$ collider.](#)
By CDF Collaboration (D. Acosta et al.).
[hep-ex/0311023].
[10.1103/PhysRevLett.93.141601.](#)
Phys.Rev.Lett. 93 (2004) 141601.

215) [The CDF calorimetry upgrade for run two \$\sqrt{s}\$.](#)
By S. Kuhlmann, Henry J. Frisch, M. Cordelli, J. Huston, R. Miller, S. Lami, R. Paoletti, N. Turini et al..
[physics/0310155].
[10.1016/j.nima.2003.10.018.](#)
Nucl.Instrum.Meth. A518 (2004) 39-41.

216) [Search for Kaluza-Klein graviton emission in \$\bar{p}p\$ collisions at \$\sqrt{s} = 1.8\$ -TeV using the missing energy signature.](#)
By CDF Collaboration (D. Acosta et al.).
[hep-ex/0309051].
[10.1103/PhysRevLett.92.121802.](#)
Phys.Rev.Lett. 92 (2004) 121802.

217) [Measurement of the average time-integrated mixing probability of \$b\$ -flavored hadrons produced at the Tevatron.](#)
By CDF Collaboration (D. Acosta et al.).
[hep-ex/0309030].

[10.1103/PhysRevD.69.012002.](#)
Phys.Rev. D69 (2004) 012002.

218) [Search for the flavor-changing neutral current decay \$D^0 \rightarrow \mu^+ \mu^-\$ in \$p\bar{p}\$ collisions at \$\sqrt{s} = 1.96\$ TeV.](#)

By CDF Collaboration (D. Acosta et al.).
[hep-ex/0308059].

[10.1103/PhysRevD.68.091101.](#)
Phys.Rev. D68 (2003) 091101.

219) [Measurement of the mass difference \$m\(D_s^+\) - m\(D^+\)\$ at {CDF} II.](#)

By CDF Collaboration (D. Acosta et al.).
[hep-ex/0310043].

[10.1103/PhysRevD.68.072004.](#)
Phys.Rev. D68 (2003) 072004.

220) [Measurement of prompt charm meson production cross sections in \$p\bar{p}\$ collisions at \$\sqrt{s} = 1.96\$ TeV.](#)

By CDF Collaboration (D. Acosta et al.).
[hep-ex/0307080].

[10.1103/PhysRevLett.91.241804.](#)
Phys.Rev.Lett. 91 (2003) 241804.

221) [Search for associated production of \$\Upsilon\$ and vector boson in \$p\bar{p}\$ collisions at \$\sqrt{s} = 1.8\$ TeV.](#)

By CDF Collaboration (D. Acosta et al.).
[10.1103/PhysRevLett.90.221803.](#)

Phys.Rev.Lett. 90 (2003) 221803.

222) [Search for lepton flavor violating decays of a heavy neutral particle in \$p\bar{p}\$ collisions at \$\sqrt{s} = 1.8\$ TeV.](#)

By CDF Collaboration (D. Acosta et al.).
[hep-ex/0307012].

[10.1103/PhysRevLett.91.171602.](#)
Phys.Rev.Lett. 91 (2003) 171602.

223) [Search for pair production of scalar top quarks in \$R\$ -parity violating decay modes in \$p\bar{p}\$ collisions at \$\sqrt{s} = 1.8\$ TeV.](#)

By CDF Collaboration (D. Acosta et al.).
[hep-ex/0305010].

[10.1103/PhysRevLett.92.051803.](#)
Phys.Rev.Lett. 92 (2004) 051803.

224) [Central pseudorapidity gaps in events with a leading antiproton at the Fermilab Tevatron \$p\bar{p}\$ collider.](#)

By CDF Collaboration (D. Acosta et al.).
[hep-ex/0303011].

[10.1103/PhysRevLett.91.011802.](#)
Phys.Rev.Lett. 91 (2003) 011802.

225) [Search for the supersymmetric partner of the top quark in dilepton events from \$p\bar{p}\$ collisions at \$\sqrt{s} = 1.8\$ TeV.](#)

By CDF Collaboration (D. Acosta et al.).
[hep-ex/0302009].

[10.1103/PhysRevLett.90.251801.](#)
Phys.Rev.Lett. 90 (2003) 251801.

- 226) [Performances of the AMS-02 electromagnetic calorimeter.](#)
By F. Cervelli, S. Di Falco, T. Lomtadze, G. Venanzoni, E. Falchini, P. Maestro, P.S. Marrocchesi, R. Paoletti et al..
- 227) [Electromagnetic calorimeter for the AMS-02 experiment.](#)
By R. Kossakowski, F. Cadoux, V. Chambert-Hermel, G. Coignet, J.M. Dubois, D. Fougeron, N. Fouque, L. Girard et al..
- 228) [The data acquisition of the MAGIC telescope.](#)
By MAGIC Collaboration (J. Cortina et al.).
- 229) [The AMS-02 electromagnetic calorimeter.](#)
By F. Cadoux, F. Cervelli, V. Chambert-Hermel, G. Chen, H. Chen, G. Coignet, S. Di Falco, J.M. Dubois et al..
[10.1016/S0920-5632\(02\)01836-4.](#)
Nucl.Phys.Proc.Suppl. 113 (2002) 159-165.
- 230) [A low-power high dynamic range front-end ASIC for imaging calorimeters.](#)
By M.G. Bagliesi, P. Maestro, P.S. Marrocchesi, M. Meucci, V. Millucci, F. Morsani, R. Paoletti, F. Pilo et al..
- 231) [Cross section for forward \$J/\psi\$ production in \$p\bar{p}\$ collisions at \$\sqrt{s} = 1.8\$ TeV.](#)
By CDF Collaboration (D. Acosta et al.).
[10.1103/PhysRevD.66.092001.](#)
Phys.Rev. D66 (2002) 092001.
- 232) [Search for long-lived charged massive particles in \$\bar{p}p\$ collisions at \$\sqrt{s} = 1.8\$ TeV.](#)
By CDF Collaboration (D. Acosta et al.).
[hep-ex/0211064].
[10.1103/PhysRevLett.90.131801.](#)
Phys.Rev.Lett. 90 (2003) 131801.
- 233) [A reduced scale e.m. calorimeter prototype for the AMS-02 experiment.](#)
By F. Cervelli, G. Chen, G. Coignet, S. Di Falco, E. Falchini, T. Lomtadze, Z. Liu, P. Maestro et al..
[10.1016/S0168-9002\(02\)00915-4.](#)
Nucl.Instrum.Meth. A490 (2002) 132-139.
- 234) [Search for a \$W'^{\pm}\$ boson decaying to a top and bottom quark pair in 1.8 TeV \$p\bar{p}\$ collisions.](#)
By CDF Collaboration (D. Acosta et al.).
[hep-ex/0209030].
[10.1103/PhysRevLett.90.081802.](#)
Phys.Rev.Lett. 90 (2003) 081802.
- 235) [Search for radiative b-hadron decays in \$p\bar{p}\$ collisions at \$\sqrt{s} = 1.8\$ TeV.](#)
By CDF Collaboration (D. Acosta et al.).
[hep-ex/0208035].
[10.1103/PhysRevD.66.112002.](#)
Phys.Rev. D66 (2002) 112002.
- 236) [Momentum distribution of charged particles in jets in dijet events in \$p\bar{p}\$ collisions at \$\sqrt{s} = 1.8\$ TeV and comparisons to perturbative QCD predictions.](#)
By CDF Collaboration (D. Acosta et al.).
[10.1103/PhysRevD.68.012003.](#)
Phys.Rev. D68 (2003) 012003.

237) [Measurement of the ratio of \$b\$ quark production cross sections in \$\bar{p}p\$ collisions at \$\sqrt{s} = 630\$ GeV and \$\sqrt{s} = 1800\$ GeV.](#)

By CDF Collaboration (D. Acosta et al.).

[hep-ex/0206019].

[10.1103/PhysRevD.66.032002.](#)

Phys.Rev. D66 (2002) 032002.

238) [Branching ratio measurements of exclusive \$B^+\$ decays to charmonium with the Collider Detector at Fermilab.](#)

By CDF Collaboration (D. Acosta et al.).

[10.1103/PhysRevD.66.052005.](#)

Phys.Rev. D66 (2002) 052005.

239) [Limits on extra dimensions and new particle production in the exclusive photon and missing energy signature in \$\bar{p}p\$ collisions at \$\sqrt{s} = 1.8\$ TeV.](#)

By CDF Collaboration (D. Acosta et al.).

[hep-ex/0205057].

[10.1103/PhysRevLett.89.281801.](#)

Phys.Rev.Lett. 89 (2002) 281801.

240) [Measurement of \$B\$ meson lifetimes using fully reconstructed \$B\$ decays produced in \$\bar{p}p\$ collisions at \$\sqrt{s} = 1.8\$ TeV.](#)

By CDF Collaboration (D. Acosta et al.).

[10.1103/PhysRevD.65.092009.](#)

Phys.Rev. D65 (2002) 092009.

241) [Search for new physics in photon lepton events in \$\bar{p}p\$ collisions at \$\sqrt{s} = 1.8\$ -TeV.](#)

By CDF Collaboration (D. Acosta et al.).

[hep-ex/0202044].

[10.1103/PhysRevLett.89.041802.](#)

Phys.Rev.Lett. 89 (2002) 041802.

242) [Comparison of the isolated direct photon cross sections in \$\bar{p}p\$ collisions at \$\sqrt{s} = 1.8\$ -TeV and \$\sqrt{s} = 0.63\$ -TeV.](#)

By CDF Collaboration (D. Acosta et al.).

[hep-ex/0201004].

[10.1103/PhysRevD.65.112003.](#)

Phys.Rev. D65 (2002) 112003.

243) [\$\Upsilon\$ production and polarization in \$\bar{p}p\$ collisions at \$\sqrt{s} = 1.8\$ -TeV.](#)

By CDF Collaboration (D. Acosta et al.).

[10.1103/PhysRevLett.88.161802.](#)

Phys.Rev.Lett. 88 (2002) 161802.

244) [Observation of the moon shadow using a new reconstruction technique in the CLUE experiment.](#)

By B. Bartoli, D. Bastieri, C. Bigongiari, R. Biral, M.A. Ciocci, M. Cresti, V. Dokuchaeva, D. Kartashov et al..

Nuovo Cim. C24 (2001) 669-674.

245) [Soft and hard interactions in \$\bar{p}p\$ collisions at \$\sqrt{s} = 1800\$ -GeV and 630-GeV.](#)

By CDF Collaboration (D. Acosta et al.).

[10.1103/PhysRevD.65.072005.](#)

Phys.Rev. D65 (2002) 072005.

246) [Measurement of the \$B^+\$ total cross section and \$B^+\$ differential cross section \$d\sigma/dp_T\$ in \$\bar{p}p\$ collisions at \$\sqrt{s} = 1.8\$ -TeV.](#)

By CDF Collaboration (D. Acosta et al.).
[hep-ph/0111359].
[10.1103/PhysRevD.65.052005](https://arxiv.org/abs/10.1103/PhysRevD.65.052005).
Phys.Rev. D65 (2002) 052005.

247) [Search for single top quark production in \$p\bar{p}\$ collisions at \$\sqrt{s}=1.8\$ -TeV.](#)
By CDF Collaboration (D. Acosta et al.).
[hep-ex/0110067].
[10.1103/PhysRevD.65.091102](https://arxiv.org/abs/10.1103/PhysRevD.65.091102).
Phys.Rev. D65 (2002) 091102.

248) [Search for the decay \$B_s \rightarrow \mu^+ \mu^- \phi\$ in \$p\bar{p}\$ collisions at \$\sqrt{s} = 1.8\$ -TeV.](#)
By CDF Collaboration (D. Acosta et al.).
[10.1103/PhysRevD.65.111101](https://arxiv.org/abs/10.1103/PhysRevD.65.111101).
Phys.Rev. D65 (2002) 111101.

249) [Charged jet evolution and the underlying event in \$p\bar{p}\$ collisions at 1.8 TeV.](#)
By CDF Collaboration (T. Affolder et al.).
[10.1103/PhysRevD.65.092002](https://arxiv.org/abs/10.1103/PhysRevD.65.092002).
Phys.Rev. D65 (2002) 092002.

250) [Search for new physics in photon lepton events in \$p\$ anti- \$p\$ collisions at \$s^{1/2} = 1.8\$ -TeV.](#)
By CDF Collaboration (D. Acosta et al.).
[hep-ex/0110015].
[10.1103/PhysRevD.66.012004](https://arxiv.org/abs/10.1103/PhysRevD.66.012004).
Phys.Rev. D66 (2002) 012004.

251) [Diffractive dijet production at \$\sqrt{s} = 630\$ GeV and 1800 GeV at the Fermilab Tevatron.](#)
By CDF Collaboration (D. Acosta et al.).
[hep-ex/0109025].
[10.1103/PhysRevLett.88.151802](https://arxiv.org/abs/10.1103/PhysRevLett.88.151802).
Phys.Rev.Lett. 88 (2002) 151802.

252) [Study of the heavy flavor content of jets produced in association with W bosons in \$p\bar{p}\$ collisions at \$\sqrt{s} = 1.8\$ TeV.](#)
By CDF Collaboration (D. Acosta et al.).
[hep-ex/0109012].
[10.1103/PhysRevD.65.052007](https://arxiv.org/abs/10.1103/PhysRevD.65.052007).
Phys.Rev. D65 (2002) 052007.

253) [A precise measurement of 180-GeV muon energy losses in iron.](#)
By ATLAS TileCal Collaboration (P. Amaral et al.).
[10.1007/s100520100645](https://arxiv.org/abs/10.1007/s100520100645).
Eur.Phys.J. C20 (2001) 487-495.

254) [Measurement of the strong coupling constant from inclusive jet production at the Tevatron \$p\bar{p}\$ collider.](#)
By CDF Collaboration (T. Affolder et al.).
[hep-ex/0108034].
[10.1103/PhysRevLett.88.042001](https://arxiv.org/abs/10.1103/PhysRevLett.88.042001).
Phys.Rev.Lett. 88 (2002) 042001.

255) [A study of \$B^0 \rightarrow J/\psi K^{\(*\)0} \pi^+ \pi^-\$ decays with the Collider Detector at Fermilab.](#)
By CDF Collaboration (T. Affolder et al.).
[hep-ex/0108022].

[10.1103/PhysRevLett.88.071801.](#)
Phys.Rev.Lett. 88 (2002) 071801.

256) [Search for new heavy particles in the \$W Z^0\$ final state in \$p\bar{p}\$ collisions at \$\sqrt{s} = 1.8\$ TeV.](#)

By CDF Collaboration (T. Affolder et al.).
[hep-ex/0108004].

[10.1103/PhysRevLett.88.071806.](#)
Phys.Rev.Lett. 88 (2002) 071806.

257) [Observation of diffractive \$J/\psi\$ production at the Fermilab Tevatron.](#)

By CDF Collaboration (T. Affolder et al.).
[hep-ex/0107071].

[10.1103/PhysRevLett.87.241802.](#)
Phys.Rev.Lett. 87 (2001) 241802.

258) [Charged particle multiplicity in jets in \$p\bar{p}\$ collisions at \$\sqrt{s} = 1.8\$ TeV.](#)

By CDF Collaboration (T. Affolder et al.).
[10.1103/PhysRevLett.87.211804.](#)

Phys.Rev.Lett. 87 (2001) 211804.

259) [Search for quark lepton compositeness and a heavy \$W'^{\prime}\$ boson using the \$e \nu\$ channel in \$p\bar{p}\$ collisions at \$\sqrt{s} = 1.8\$ TeV.](#)

By CDF Collaboration (T. Affolder et al.).
[hep-ex/0107008].

[10.1103/PhysRevLett.87.231803.](#)
Phys.Rev.Lett. 87 (2001) 231803.

260) [Searches for new physics in events with a photon and b-quark jet at CDF.](#)

By CDF Collaboration (T. Affolder et al.).
[hep-ex/0106012].

[10.1103/PhysRevD.65.052006.](#)
Phys.Rev. D65 (2002) 052006.

261) [A two-level pattern trigger for the MAGIC telescope.](#)

By D. Bastieri, M. Bettini, C. Bigongiari, D. Corti, R. Isocrate, M. Mariotti, L. Peruzzo, A. Saggion et al..
[10.1016/S0168-9002\(00\)01289-4.](#)

Nucl.Instrum.Meth. A461 (2001) 521-523.

262) [Shower reconstruction in the CLUE experiment.](#)

By B. Bartoli, D. Bastieri, C. Bigongiari, M.A. Ciocci, D. Cosulich, M. Cresti, V. Dokuchaeva, D. Kartashov et al..

[10.1016/S0168-9002\(00\)01253-5.](#)

Nucl.Instrum.Meth. A461 (2001) 311-313.

263) [Search for gluinos and squarks using like-sign dileptons in \$p\bar{p}\$ collisions at \$\sqrt{s} = 1.8\$ TeV.](#)

By CDF Collaboration (T. Affolder et al.).
[hep-ex/0106061].

[10.1103/PhysRevLett.87.251803.](#)
Phys.Rev.Lett. 87 (2001) 251803.

264) [Measurement of \$d\(\sigma\)/dM\$ and forward-backward charge asymmetry for high mass Drell-Yan \$e^+e^-\$ pairs from \$p\bar{p}\$ collisions at \$\sqrt{s} = 1.8\$ TeV.](#)

By CDF Collaboration (T. Affolder et al.).
[hep-ex/0106047].

[10.1103/PhysRevLett.87.131802](#).
Phys.Rev.Lett. 87 (2001) 131802.

265) [Observation of gamma-sources using a new reconstruction technique in the CLUE experiment.](#)
By CLUE Collaboration (B. Bartoli et al.).
[10.1016/S0920-5632\(01\)01266-X](#).
Nucl.Phys.Proc.Suppl. 97 (2001) 211-214.

266) [Cross section and heavy quark composition of \$\gamma + \mu\$ events produced in \$p\bar{p}\$ collisions.](#)
By CDF Collaboration (T. Affolder et al.).
[hep-ex/0106004].
[10.1103/PhysRevD.65.012003](#).
Phys.Rev. D65 (2002) 012003.

267) [Search for gluinos and scalar quarks in \$p\bar{p}\$ collisions at \$\sqrt{s} = 1.8\$ TeV using the missing energy plus multijets signature.](#)
By CDF Collaboration (T. Affolder et al.).
[hep-ex/0106001].
[10.1103/PhysRevLett.88.041801](#).
Phys.Rev.Lett. 88 (2002) 041801.

268) [Production of \$\chi_{c1}\$ and \$\chi_{c2}\$ in \$p\bar{p}\$ collisions at \$\sqrt{s} = 1.8\$ TeV.](#)
By CDF Collaboration (T. Affolder et al.).
[10.1103/PhysRevLett.86.3963](#).
Phys.Rev.Lett. 86 (2001) 3963-3968.

269) [Double diffraction dissociation at the Fermilab Tevatron collider.](#)
By CDF Collaboration (T. Affolder et al.).
[hep-ex/0107070].
[10.1103/PhysRevLett.87.141802](#).
Phys.Rev.Lett. 87 (2001) 141802.

270) [Search for narrow diphoton resonances and for \$\gamma\gamma + W/Z\$ signatures in \$p\bar{p}\$ collisions at \$\sqrt{s} = 1.8\$ TeV.](#)
By CDF Collaboration (T. Affolder et al.).
[hep-ex/0105066].
[10.1103/PhysRevD.64.092002](#).
Phys.Rev. D64 (2001) 092002.

271) [Hadron energy reconstruction for the ATLAS calorimetry in the framework of the nonparametrical method.](#)
By ATLAS Collaboration (S. Akhmadalev et al.).
[hep-ex/0104002].
[10.1016/S0168-9002\(01\)01229-3](#).
Nucl.Instrum.Meth. A480 (2002) 508-523.

272) [Measurement of the inclusive jet cross section in \$p\bar{p}\$ collisions at \$\sqrt{s} = 1.8\$ TeV.](#)
By CDF Collaboration (T. Affolder et al.).
[hep-ph/0102074].
[10.1103/PhysRevD.65.039903](#), [10.1103/PhysRevD.64.032001](#).
Phys.Rev. D64 (2001) 032001, Erratum-ibid. D65 (2002) 039903.

273) [Measurement of the \$t\bar{t}\$ production cross section in \$p\bar{p}\$ collisions at \$\sqrt{s} = 1.8\$ TeV.](#)
By CDF Collaboration (T. Affolder et al.).

[hep-ex/0101036].

[10.1103/PhysRevD.67.119901](#), [10.1103/PhysRevD.64.032002](#).

Phys.Rev. D64 (2001) 032002, Erratum-ibid. D67 (2003) 119901.

274) [The CLUE telescope array: Observation of gamma-ray sources.](#)

By CLUE Collaboration (B. Bartoli et al.).

275) [First measurement of the ratio \$B\(t \rightarrow W b\)/B\(t \rightarrow W q\)\$ and associated limit on the CKM element \$|V_{tb}|\$.](#)

By CDF Collaboration (T. Affolder et al.).

[hep-ex/0012029].

[10.1103/PhysRevLett.86.3233](#).

Phys.Rev.Lett. 86 (2001) 3233-3238.

276) [Measurement of the two-jet differential cross section in \$p\bar{p}\$ collisions at \$\sqrt{s} = 1800\$ GeV.](#)

By CDF Collaboration (T. Affolder et al.).

[hep-ex/0012013].

[10.1103/PhysRevD.65.039902](#), [10.1103/PhysRevD.64.012001](#).

Phys.Rev. D64 (2001) 012001, Erratum-ibid. D65 (2002) 039902.

277) [Measurement of the top quark \$p_T\$ distribution.](#)

By CDF Collaboration (T. Affolder et al.).

[10.1103/PhysRevLett.87.102001](#).

Phys.Rev.Lett. 87 (2001) 102001.

278) [Search for the supersymmetric partner of the top quark in \$p\bar{p}\$ collisions at \$\sqrt{s} = 1.8\$ TeV.](#)

By CDF Collaboration (T. Affolder et al.).

[hep-ex/0011004].

[10.1103/PhysRevD.63.091101](#).

Phys.Rev. D63 (2001) 091101.

279) [Search for neutral supersymmetric Higgs bosons in \$p\bar{p}\$ collisions at \$\sqrt{s} = 1.8\$ TeV.](#)

By CDF Collaboration (T. Affolder et al.).

[hep-ex/0010052].

[10.1103/PhysRevLett.86.4472](#).

Phys.Rev.Lett. 86 (2001) 4472-4478.

280) [Test of enhanced leading order QCD in \$W\$ boson plus jets events from 1.8 TeV \$p\bar{p}\$ collisions.](#)

By CDF Collaboration (T. Affolder et al.).

[10.1103/PhysRevD.63.072003](#).

Phys.Rev. D63 (2001) 072003.

281) [Results from a new combined test of an electromagnetic liquid argon calorimeter with a hadronic scintillating-tile calorimeter.](#)

By ATLAS Collaboration (S. Akhmadalev et al.).

[10.1016/S0168-9002\(00\)00153-4](#).

Nucl.Instrum.Meth. A449 (2000) 461-477.

282) [Measurement of the \$W\$ boson mass with the Collider Detector at Fermilab.](#)

By CDF Collaboration (T. Affolder et al.).

[hep-ex/0007044].

[10.1103/PhysRevD.64.052001](#).

Phys.Rev. D64 (2001) 052001.

283) [Measurement of the decay amplitudes of \$B^0 \rightarrow J/\psi K^{*0}\$ and \$B^0 \rightarrow J/\psi \phi\$ decays.](#)

By CDF Collaboration (T. Affolder et al.).

[hep-ex/0007034].

[10.1103/PhysRevLett.85.4668.](#)

Phys.Rev.Lett. 85 (2000) 4668-4673.

284) [Measurement of the top quark mass with the Collider Detector at Fermilab.](#)

By CDF Collaboration (T. Affolder et al.).

[hep-ex/0006028].

[10.1103/PhysRevD.63.032003.](#)

Phys.Rev. D63 (2001) 032003.

285) [Measurement of \$d\(\sigma\)/dy\$ for high mass Drell-Yan \$e^+e^-\$ pairs from \$p\bar{p}\$ collisions at \$\sqrt{s} = 1.8\$ TeV.](#)

By CDF Collaboration (T. Affolder et al.).

[hep-ex/0006025].

[10.1103/PhysRevD.63.011101.](#)

Phys.Rev. D63 (2001) 011101.

286) [Dijet production by double pomeron exchange at the Fermilab Tevatron.](#)

By CDF Collaboration (T. Affolder et al.).

[10.1103/PhysRevLett.85.4215.](#)

Phys.Rev.Lett. 85 (2000) 4215-4220.

287) [Measurement of \$J/\psi\$ and \$\psi\(2S\)\$ polarization in \$p\bar{p}\$ collisions at \$\sqrt{s} = 1.8\$ TeV.](#)

By CDF Collaboration (T. Affolder et al.).

[hep-ex/0004027].

[10.1103/PhysRevLett.85.2886.](#)

Phys.Rev.Lett. 85 (2000) 2886-2891.

288) [Direct measurement of the \$W\$ boson width in \$p\bar{p}\$ collisions at \$\sqrt{s} = 1.8\$ TeV.](#)

By CDF Collaboration (T. Affolder et al.).

[hep-ex/0004017].

[10.1103/PhysRevLett.85.3347.](#)

Phys.Rev.Lett. 85 (2000) 3347-3352.

289) [Search for second and third generation leptoquarks including production via technicolor interactions in \$p\bar{p}\$ collisions at \$\sqrt{s} = 1.8\$ TeV.](#)

By CDF Collaboration (T. Affolder et al.).

[hep-ex/0004003].

[10.1103/PhysRevLett.85.2056.](#)

Phys.Rev.Lett. 85 (2000) 2056-2061.

290) [Limits on gravitino production and new processes with large missing transverse energy in \$p\bar{p}\$ collisions at \$\sqrt{s} = 1.8\$ TeV.](#)

By CDF Collaboration (T. Affolder et al.).

[hep-ex/0003026].

[10.1103/PhysRevLett.85.1378.](#)

Phys.Rev.Lett. 85 (2000) 1378-1383.

291) [Diffractive dijets with a leading antiproton in \$p\bar{p}\$ collisions at \$\sqrt{s} = 1800\$ GeV.](#)

By CDF Collaboration (T. Affolder et al.).

[10.1103/PhysRevLett.84.5043.](#)

Phys.Rev.Lett. 84 (2000) 5043-5048.

292) [Search for new particles decaying to \$t\bar{t}\$ in \$p\bar{p}\$ collisions at \$\sqrt{s} = 1.8\$ TeV.](#)

By CDF Collaboration (T. Affolder et al.).

[hep-ex/0003005].

[10.1103/PhysRevLett.85.2062.](#)

Phys.Rev.Lett. 85 (2000) 2062-2067.

293) [The CLUE experiment running with 8 telescopes: Observations of gamma sources and runs on Moon.](#)

By D. Bastieri, B. Bartoli, C. Bigongiari, R. Biral, M.A. Ciocci, D. Cosulich, M. Cresti, D. Kartashov et al..

294) [A measurement of the differential dijet mass cross section in \$p\bar{p}\$ collisions at \$\sqrt{s} = 1.8\$ TeV.](#)

By CDF Collaboration (T. Affolder et al.).

[hep-ex/9912022].

[10.1103/PhysRevD.61.091101.](#)

Phys.Rev. D61 (2000) 091101.

295) [Search for scalar top quark production in \$p\bar{p}\$ collisions at \$\sqrt{s} = 1.8\$ TeV.](#)

By CDF Collaboration (T. Affolder et al.).

[hep-ex/9912018, hep-ex/9912018].

[10.1103/PhysRevLett.84.5273.](#)

Phys.Rev.Lett. 84 (2000) 5273-5278.

296) [Observation of orbitally excited \$B\$ mesons in \$p\bar{p}\$ collisions at \$\sqrt{s} = 1.8\$ TeV.](#)

By CDF Collaboration (T. Affolder et al.).

[10.1103/PhysRevD.64.072002.](#)

Phys.Rev. D64 (2001) 072002.

297) [Search for the charged Higgs boson in the decays of top quark pairs in the \$e\tau\$ and \$\mu\tau\$ channels at \$\sqrt{s} = 1.8\$ TeV.](#)

By CDF Collaboration (T. Affolder et al.).

[hep-ex/9912013].

[10.1103/PhysRevD.62.012004.](#)

Phys.Rev. D62 (2000) 012004.

298) [Search for scalar top and scalar bottom quarks in \$p\bar{p}\$ collisions at \$\sqrt{s} = 1.8\$ TeV.](#)

By CDF Collaboration (T. Affolder et al.).

[hep-ex/9910049].

[10.1103/PhysRevLett.84.5704.](#)

Phys.Rev.Lett. 84 (2000) 5704-5709.

299) [Production of \$\Upsilon\(1S\)\$ mesons from \$\chi_b\$ decays in \$p\bar{p}\$ collisions at \$\sqrt{s} = 1.8\$ TeV.](#)

By CDF Collaboration (T. Affolder et al.).

[hep-ex/9910025].

[10.1103/PhysRevLett.84.2094.](#)

Phys.Rev.Lett. 84 (2000) 2094-2099.

300) [Measurement of \$b\$ quark fragmentation fractions in the production of strange and light \$B\$ mesons in \$p\bar{p}\$ collisions at \$\sqrt{s} = 1.8\$ TeV.](#)

By CDF Collaboration (F. Abe et al.).

[10.1103/PhysRevD.60.092005.](#)

Phys.Rev. D60 (1999) 092005.

301) [Search for a \$W^{\prime}\$ boson via the decay mode \$W^{\prime} \rightarrow \mu \nu_{\mu}\$ in 1.8 TeV \$p\bar{p}\$ collisions.](#)

By CDF Collaboration (F. Abe et al.).

[hep-ex/9910004].

[10.1103/PhysRevLett.84.5716.](#)

Phys.Rev.Lett. 84 (2000) 5716-5721.

302) [Measurement of the helicity of \$W\$ bosons in top quark decays.](#)

By CDF Collaboration (T. Affolder et al.).

[hep-ex/9909042].

[10.1103/PhysRevLett.84.216.](#)

Phys.Rev.Lett. 84 (2000) 216-221.

303) [Search for a fourth-generation quark more massive than the \$Z_0\$ boson in \$p\bar{p}\$ collisions at \$\sqrt{s} = 1.8\$ TeV.](#)

By CDF Collaboration (T. Affolder et al.).

[hep-ex/9909027].

[10.1103/PhysRevLett.84.835.](#)

Phys.Rev.Lett. 84 (2000) 835-840.

304) [Measurement of \$b\$ quark fragmentation fractions in \$p\bar{p}\$ collisions at \$\sqrt{s} = 1.8\$ TeV.](#)

By CDF Collaboration (T. Affolder et al.).

[hep-ex/9909011].

[10.1103/PhysRevLett.84.1663.](#)

Phys.Rev.Lett. 84 (2000) 1663-1668.

305) [A measurement of \$\sin\(2\beta\)\$ from \$B \rightarrow J/\psi K^0 S\$ with the {CDF} detector.](#)

By CDF Collaboration (T. Affolder et al.).

[hep-ex/9909003].

[10.1103/PhysRevD.61.072005.](#)

Phys.Rev. D61 (2000) 072005.

306) [The transverse momentum and total cross section of \$e^+e^-\$ pairs in the \$Z\$ boson region from \$p\bar{p}\$ collisions at \$\sqrt{s} = 1.8\$ TeV.](#)

By CDF Collaboration (T. Affolder et al.).

[hep-ex/0001021].

[10.1103/PhysRevLett.84.845.](#)

Phys.Rev.Lett. 84 (2000) 845-850.

307) [Observation of diffractive beauty production at the Fermilab Tevatron.](#)

By CDF Collaboration (T. Affolder et al.).

[10.1103/PhysRevLett.84.232.](#)

Phys.Rev.Lett. 84 (2000) 232-237.

308) [Reconstruction of showers at TeV energy by the CLUE experiment and its application to recent data.](#)

By B. Bartoli, D. Bastieri, M. Cresti, M. Mariotti, L. Peruzzo, R. Sacco, A. Saggion, G. Sartori et al..

Nucl.Phys.Proc.Suppl. 75A (1999) 373-376.

309) [Measurement of the \$B^0 \rightarrow \bar{B}^0\$ oscillation frequency using \$e^+e^- \rightarrow D^{*+}\$ pairs and lepton flavor tags.](#)

By CDF Collaboration (T. Affolder et al.).

[hep-ex/9907053].

[10.1103/PhysRevD.60.112004.](#)

Phys.Rev. D60 (1999) 112004.

- 310) [Search for color singlet technicolor particles in \$p\bar{p}\$ collisions at \$\sqrt{s} = 1.8\$ TeV.](#)
 By CDF Collaboration (T. Affolder et al.).
[10.1103/PhysRevLett.84.1110](#).
 Phys.Rev.Lett. 84 (2000) 1110-1115.
- 311) [Search for the flavor-changing neutral current decays \$B^+ \rightarrow \mu^+ \mu^- K^+\$ and \$B^0 \rightarrow \mu^+ \mu^- K^{*0}\$.](#)
 By CDF Collaboration (T. Affolder et al.).
 [hep-ex/9905004].
[10.1103/PhysRevLett.83.3378](#).
 Phys.Rev.Lett. 83 (1999) 3378-3383.
- 312) [Hadronic shower development in iron scintillator tile calorimetry.](#)
 By ATLAS/Tilecal Collaboration (P. Amaral et al.).
 [hep-ex/9904032].
[10.1016/S0168-9002\(99\)01020-7](#).
 Nucl.Instrum.Meth. A443 (2000) 51-70.
- 313) [Measurement of the \$B_d^0 \rightarrow B_d^0\$ oscillation frequency using dimuon data in \$p\bar{p}\$ collisions at \$\sqrt{s} = 1.8\$ TeV.](#)
 By CDF Collaboration (F. Abe et al.).
[10.1103/PhysRevD.60.051101](#).
 Phys.Rev. D60 (1999) 051101.
- 314) [Measurement of \$B^0 \rightarrow B^0\$ flavor oscillations using jet-charge and lepton flavor tagging in \$p\bar{p}\$ collisions at \$\sqrt{s} = 1.8\$ TeV.](#)
 By CDF Collaboration (F. Abe et al.).
 [hep-ex/9903011].
[10.1103/PhysRevD.60.072003](#).
 Phys.Rev. D60 (1999) 072003.
- 315) [Measurement of the associated \$\gamma + \mu^\pm\$ production cross section in \$p\bar{p}\$ collisions at \$\sqrt{s} = 1.8\$ TeV.](#)
 By CDF Collaboration (F. Abe et al.).
 [hep-ex/9902001].
[10.1103/PhysRevD.60.092003](#).
 Phys.Rev. D60 (1999) 092003.
- 316) [Preliminary results from the CLUE Cherenkov UV experiment.](#)
 By B. Bartoli, D. Bastieri, M. Cresti, M. Mariotti, L. Peruzzo, R. Sacco, A. Saggion, G. Sartori et al..
- 317) [A search for \$B_s^0 \rightarrow B_s^0\$ oscillations using the semileptonic decay \$B_s^0 \rightarrow \phi \ell \ell^+ X \nu\$.](#)
 By CDF Collaboration (F. Abe et al.).
[10.1103/PhysRevLett.82.3576](#).
 Phys.Rev.Lett. 82 (1999) 3576-3580.
- 318) [Measurement of \$b\bar{b}\$ rapidity correlations in \$p\bar{p}\$ collisions at \$\sqrt{s} = 1.8\$ TeV.](#)
 By CDF Collaboration (F. Abe et al.).
[10.1103/PhysRevD.61.032001](#).
 Phys.Rev. D61 (2000) 032001.
- 319) [Search for \$R\$ -parity violating supersymmetry using like-sign dielectrons in \$p\bar{p}\$ collisions at \$\sqrt{s} = 1.8\$ TeV.](#)
 By CDF Collaboration (F. Abe et al.).

[hep-ex/9908063].

[10.1103/PhysRevLett.83.2133](https://arxiv.org/abs/10.1103/PhysRevLett.83.2133).

Phys.Rev.Lett. 83 (1999) 2133-2138.

320) [Events with a rapidity gap between jets in \$\bar{p}p\$ collisions at \$\sqrt{s} = 630\$ GeV.](#)

By CDF Collaboration (F. Abe et al.).

[10.1103/PhysRevLett.81.5278](https://arxiv.org/abs/10.1103/PhysRevLett.81.5278).

Phys.Rev.Lett. 81 (1998) 5278-5283.

321) [Search for third-generation leptoquarks from technicolor models in \$p\bar{p}\$ collisions at \$\sqrt{s} = 1.8\$ TeV.](#)

By CDF Collaboration (F. Abe et al.).

[10.1103/PhysRevLett.82.3206](https://arxiv.org/abs/10.1103/PhysRevLett.82.3206).

Phys.Rev.Lett. 82 (1999) 3206.

322) [Kinematics of \$t\bar{t}\$ events at CDF.](#)

By CDF Collaboration (F. Abe et al.).

[10.1103/PhysRevD.59.092001](https://arxiv.org/abs/10.1103/PhysRevD.59.092001).

Phys.Rev. D59 (1999) 092001.

323) [Search for a technicolor \$\omega\(T\)\$ particle in events with a photon and a b quark jet at CDF.](#)

By CDF Collaboration (F. Abe et al.).

[hep-ex/9810031].

[10.1103/PhysRevLett.83.3124](https://arxiv.org/abs/10.1103/PhysRevLett.83.3124).

Phys.Rev.Lett. 83 (1999) 3124-3129.

324) [Measurement of the top quark mass with the collider detector at Fermilab.](#)

By CDF Collaboration (F. Abe et al.).

[hep-ex/9810029].

[10.1103/PhysRevLett.82.271](https://arxiv.org/abs/10.1103/PhysRevLett.82.271).

Phys.Rev.Lett. 82 (1999) 271-276, Erratum-ibid. 82 (1999) 2808-2809.

325) [The CLUE trigger: A VME based two level trigger for VHE experiments.](#)

By D. Alexandreas, B. Bartoli, D. Bastieri, R. Biral, C. Bigongiari, G. Chiarelli, R. Cappuccio, E. Cocca et al..

[10.1016/S0168-9002\(97\)01350-8](https://arxiv.org/abs/10.1016/S0168-9002(97)01350-8).

Nucl.Instrum.Meth. A409 (1998) 679-681.

326) [The CLUE experiment operating with 4 telescopes in La Palma.](#)

By D. Alexandreas, D. Bastieri, M. Cresti, M. Mariotti, L. Peruzzo, R. Sacco, A. Saggion, G. Sartori et al..

[10.1016/S0168-9002\(98\)00133-8](https://arxiv.org/abs/10.1016/S0168-9002(98)00133-8).

Nucl.Instrum.Meth. A409 (1998) 488-491.

327) [Measurement of \$Z^0\$ and Drell-Yan production cross section using dimuons in \$\bar{p}p\$ collisions at \$\sqrt{s} = 1.8\$ TeV.](#)

By CDF Collaboration (F. Abe et al.).

[10.1103/PhysRevD.59.052002](https://arxiv.org/abs/10.1103/PhysRevD.59.052002).

Phys.Rev. D59 (1999) 052002.

328) [Search for new particles decaying to \$b\bar{b}\$ in \$p\bar{p}\$ collisions at \$\sqrt{s} = 1.8\$ TeV.](#)

By CDF Collaboration (F. Abe et al.).

[hep-ex/9809022].

[10.1103/PhysRevLett.82.2038](https://arxiv.org/abs/10.1103/PhysRevLett.82.2038).

Phys.Rev.Lett. 82 (1999) 2038-2043.

329) [Search for the decays \$B_s^0 \rightarrow B_d^0 \ell^+ \ell^-\$ and Pati-Salam leptoquarks.](#)

By CDF Collaboration (F. Abe et al.).
[10.1103/PhysRevLett.81.5742](#).
Phys.Rev.Lett. 81 (1998) 5742-5747.

330) [Measurement of the lepton charge asymmetry in \$W\$ boson decays produced in \$p\bar{p}\$ collisions.](#)

By CDF Collaboration (F. Abe et al.).
[hep-ex/9809001].
[10.1103/PhysRevLett.81.5754](#).
Phys.Rev.Lett. 81 (1998) 5754-5759.

331) [Search for Higgs bosons produced in association with a vector boson in \$p\bar{p}\$ collisions at \$\sqrt{s} = 1.8\$ TeV.](#)

By CDF Collaboration (F. Abe et al.).
[10.1103/PhysRevLett.81.5748](#).
Phys.Rev.Lett. 81 (1998) 5748-5753.

332) [Measurement of the \$B_s^0\$ meson lifetime using semileptonic decays.](#)

By CDF Collaboration (F. Abe et al.).
[hep-ex/9808003].
[10.1103/PhysRevD.59.032004](#).
Phys.Rev. D59 (1999) 032004.

333) [Search for second generation leptoquarks in the dimuon plus dijet channel of \$p\bar{p}\$ collisions at \$\sqrt{s} = 1.8\$ TeV.](#)

By CDF Collaboration (F. Abe et al.).
[hep-ex/9810024].
[10.1103/PhysRevLett.81.4806](#).
Phys.Rev.Lett. 81 (1998) 4806-4811.

334) [Searches for new physics in diphoton events in \$p\bar{p}\$ collisions at \$\sqrt{s} = 1.8\$ TeV.](#)

By CDF Collaboration (F. Abe et al.).
[hep-ex/9806034].
[10.1103/PhysRevD.59.092002](#).
Phys.Rev. D59 (1999) 092002.

335) [Measurement of the \$B_{d^0 - \overline{B}^0}\$ flavor oscillation frequency and study of same side flavor tagging of \$B\$ mesons in \$p\bar{p}\$ collisions.](#)

By CDF Collaboration (F. Abe et al.).
[hep-ex/9806026].
[10.1103/PhysRevD.59.032001](#).
Phys.Rev. D59 (1999) 032001.

336) [Measurement of the CP-violation parameter \$\sin\(2\beta\)\$ in \$B_{d^0} / \overline{B}^0 \rightarrow J/\psi K^0 S\$ decays.](#)

By CDF Collaboration (F. Abe et al.).
[hep-ex/9806025].
[10.1103/PhysRevLett.81.5513](#).
Phys.Rev.Lett. 81 (1998) 5513-5518.

337) [Improved measurement of the \$B^+\$ and \$\overline{B}^0\$ meson lifetimes using semileptonic decays.](#)

By CDF Collaboration (F. Abe et al.).
[hep-ex/9806018].
[10.1103/PhysRevD.58.092002](#).
Phys.Rev. D58 (1998) 092002.

- 338) [Observation of the \$B_c\$ meson in \$p\bar{p}\$ collisions at \$\sqrt{s} = 1.8\$ TeV.](#)
 By CDF Collaboration (F. Abe et al.).
 [hep-ex/9805034].
[10.1103/PhysRevLett.81.2432](#).
 Phys.Rev.Lett. 81 (1998) 2432-2437.
- 339) [Search for long-lived parents of \$Z^0\$ bosons in \$p\bar{p}\$ collisions at \$\sqrt{s} = 1.8\$ TeV.](#)
 By CDF Collaboration (F. Abe et al.).
 [hep-ex/9805017].
[10.1103/PhysRevD.58.051102](#).
 Phys.Rev. D58 (1998) 051102.
- 340) [Measurement of the \$\frac{\sigma\(W + \geq 1 \text{ jet}\)}{\sigma\(W\)}\$ cross section ratio from \$p\bar{p}\$ collisions at \$\sqrt{s} = 1.8\$ TeV.](#)
 By CDF Collaboration (F. Abe et al.).
[10.1103/PhysRevLett.81.1367](#).
 Phys.Rev.Lett. 81 (1998) 1367-1372.
- 341) [Observation of \$B_c\$ mesons in \$p\bar{p}\$ collisions at \$\sqrt{s} = 1.8\$ TeV.](#)
 By CDF Collaboration (F. Abe et al.).
 [hep-ex/9804014].
[10.1103/PhysRevD.58.112004](#).
 Phys.Rev. D58 (1998) 112004.
- 342) [Search for the rare decay \$W^\pm \rightarrow D_s^{+-} \gamma\$ in \$p\bar{p}\$ collisions at \$\sqrt{s} = 1.8\$ TeV.](#)
 By CDF Collaboration (F. Abe et al.).
[10.1103/PhysRevD.58.091101](#).
 Phys.Rev. D58 (1998) 091101.
- 343) [Search for the rare decay \$W^\pm \rightarrow \pi^\pm + \gamma\$ in \$p\bar{p}\$ collisions at \$\sqrt{s} = 1.8\$ TeV.](#)
 By CDF Collaboration (F. Abe et al.).
[10.1103/PhysRevD.58.031101](#).
 Phys.Rev. D58 (1998) 031101.
- 344) [Observation of \$B^+ \rightarrow \psi\(2S\) K^+\$ and \$B^0 \rightarrow \psi\(2S\) K^0\(892\)\$ decays and measurements of \$B\$ meson branching fractions into \$J/\psi\$ and \$\psi\(2S\)\$ final states.](#)
 By CDF Collaboration (F. Abe et al.).
 [hep-ex/9803013].
[10.1103/PhysRevD.58.072001](#).
 Phys.Rev. D58 (1998) 072001.
- 345) [Search for chargino - neutralino associated production at the Fermilab Tevatron collider.](#)
 By CDF Collaboration (F. Abe et al.).
 [hep-ex/9803015].
[10.1103/PhysRevLett.80.5275](#).
 Phys.Rev.Lett. 80 (1998) 5275-5280.
- 346) [Searches for new physics in diphoton events in \$p\bar{p}\$ collisions at \$\sqrt{s} = 1.8\$ TeV.](#)
 By CDF Collaboration (F. Abe et al.).
 [hep-ex/9801019].
[10.1103/PhysRevLett.81.1791](#).
 Phys.Rev.Lett. 81 (1998) 1791-1796.

- 347) [Search for the decays \$B^0 \rightarrow \mu^+ \mu^-\$ and \$B^s \rightarrow \mu^+ \mu^-\$ in \$p\bar{p}\$ collisions at \$\sqrt{s} = 1.8\$ TeV.](#)
By CDF Collaboration (F. Abe et al.).
[10.1103/PhysRevD.57.R3811.](#)
Phys.Rev. D57 (1998) 3811-3816.
- 348) [Jet pseudorapidity distribution in direct photon events in \$p\bar{p}\$ collisions at \$\sqrt{s} = 1.8\$ TeV.](#)
By CDF Collaboration (F. Abe et al.).
[10.1103/PhysRevD.57.1359.](#)
Phys.Rev. D57 (1998) 1359-1365.
- 349) [Observation of hadronic W decays in \$t\bar{t}\$ events with the Collider Detector at Fermilab.](#)
By CDF Collaboration (F. Abe et al.).
[hep-ex/9711004].
[10.1103/PhysRevLett.80.5720.](#)
Phys.Rev.Lett. 80 (1998) 5720-5725.
- 350) [Measurement of \$B\$ hadron lifetimes using \$J/\psi\$ final states at CDF.](#)
By CDF Collaboration (F. Abe et al.).
[10.1103/PhysRevD.57.5382.](#)
Phys.Rev. D57 (1998) 5382-5401.
- 351) [Measurement of the top quark mass and \$t\bar{t}\$ production cross section from dilepton events at the Collider Detector at Fermilab.](#)
By CDF Collaboration (F. Abe et al.).
[hep-ex/9802017].
[10.1103/PhysRevLett.80.2779.](#)
Phys.Rev.Lett. 80 (1998) 2779-2784.
- 352) [Measurement of the top quark mass.](#)
By CDF Collaboration (F. Abe et al.).
[hep-ex/9801014].
[10.1103/PhysRevLett.80.2767.](#)
Phys.Rev.Lett. 80 (1998) 2767-2772.
- 353) [Measurement of the \$t\bar{t}\$ production cross section in \$p\bar{p}\$ collisions at \$\sqrt{s} = 1.8\$ TeV.](#)
By CDF Collaboration (F. Abe et al.).
[hep-ex/9710008].
[10.1103/PhysRevLett.80.2773.](#)
Phys.Rev.Lett. 80 (1998) 2773-2778.
- 354) [Measurement of the \$B^0 \rightarrow \bar{B}^0\$ oscillation frequency using \$\pi B\$ meson charge-flavor correlations in \$p\bar{p}\$ collisions at \$\sqrt{s} = 1.8\$ TeV.](#)
By CDF Collaboration (F. Abe et al.).
[hep-ex/9712004].
[10.1103/PhysRevLett.80.2057.](#)
Phys.Rev.Lett. 80 (1998) 2057-2062.
- 355) [Search for flavor-changing neutral current decays of the top quark in \$p\bar{p}\$ collisions at \$\sqrt{s} = 1.8\$ TeV.](#)
By CDF Collaboration (F. Abe et al.).
[10.1103/PhysRevLett.80.2525.](#)
Phys.Rev.Lett. 80 (1998) 2525-2530.

- 356) [Properties of jets in \$W\$ boson events from 1.8 TeV \$\bar{p}p\$ collisions.](#)
By CDF Collaboration (F. Abe et al.).
[hep-ex/9709016].
[10.1103/PhysRevLett.79.4760](#).
Phys.Rev.Lett. 79 (1997) 4760-4765.
- 357) [Measurement of the differential cross section for events with large total transverse energy in \$p\bar{p}\$ collisions at \$\sqrt{s} = 1.8\$ TeV.](#)
By CDF Collaboration (F. Abe et al.).
[10.1103/PhysRevLett.80.3461](#).
Phys.Rev.Lett. 80 (1998) 3461-3466.
- 358) [Dijet production by color - singlet exchange at the Fermilab Tevatron.](#)
By CDF Collaboration (F. Abe et al.).
[10.1103/PhysRevLett.80.1156](#).
Phys.Rev.Lett. 80 (1998) 1156-1161.
- 359) [Search for first generation leptoquark pair production in \$p\bar{p}\$ collisions at \$\sqrt{s} = 1.8\$ TeV.](#)
By CDF Collaboration (F. Abe et al.).
[hep-ex/9708017].
[10.1103/PhysRevLett.79.4327](#).
Phys.Rev.Lett. 79 (1997) 4327-4332.
- 360) [Search for new particles decaying into \$b\bar{b}\$ and produced in association with \$W\$ bosons decaying into \$e\nu\$ or \$\mu\nu\$ at the Tevatron.](#)
By CDF Collaboration (F. Abe et al.).
[10.1103/PhysRevLett.79.3819](#).
Phys.Rev.Lett. 79 (1997) 3819-3824.
- 361) [Limits on quark-lepton compositeness scales from dileptons produced in 1.8 TeV \$p\bar{p}\$ collisions.](#)
By CDF Collaboration (F. Abe et al.).
[10.1103/PhysRevLett.79.2198](#).
Phys.Rev.Lett. 79 (1997) 2198-2203.
- 362) [Search for new gauge bosons decaying into dileptons in \$\bar{p}p\$ collisions at \$\sqrt{s} = 1.8\$ TeV.](#)
By CDF Collaboration (F. Abe et al.).
[10.1103/PhysRevLett.79.2192](#).
Phys.Rev.Lett. 79 (1997) 2192-2197.
- 363) [Properties of photon plus two-jet events in \$\bar{p}p\$ collisions at \$\sqrt{s} = 1.8\$ TeV.](#)
By CDF Collaboration (F. Abe et al.).
[10.1103/PhysRevD.57.67](#).
Phys.Rev. D57 (1998) 67-77.
- 364) [Properties of six-jet events with large six-jet mass at the Fermilab \$p\bar{p}\$ collider.](#)
By CDF Collaboration (F. Abe et al.).
[10.1103/PhysRevD.56.2532](#).
Phys.Rev. D56 (1997) 2532-2543.
- 365) [Double parton scattering in \$\bar{p}p\$ collisions at \$\sqrt{s} = 1.8\$ TeV.](#)
By CDF Collaboration (F. Abe et al.).
[10.1103/PhysRevD.56.3811](#).
Phys.Rev. D56 (1997) 3811-3832.

366) [The mu tau and e tau decays of top quark pairs produced in \$p\bar{p}\$ collisions at \$\sqrt{s} = 1.8\$ TeV.](#)

By CDF Collaboration (F. Abe et al.).
[hep-ex/9704007].

[10.1103/PhysRevLett.79.3585.](#)

Phys.Rev.Lett. 79 (1997) 3585-3590.

367) [Measurement of double parton scattering in \$p\bar{p}\$ collisions at \$\sqrt{s} = 1.8\$ TeV.](#)

By CDF Collaboration (F. Abe et al.).

[10.1103/PhysRevLett.79.584.](#)

Phys.Rev.Lett. 79 (1997) 584-589.

368) [Search for charged Higgs decays of the top quark using hadronic decays of the tau lepton.](#)

By CDF Collaboration (F. Abe et al.).

[hep-ex/9704003].

[10.1103/PhysRevLett.79.357.](#)

Phys.Rev.Lett. 79 (1997) 357-362.

369) [First observation of the all hadronic decay of \$t\bar{t}\$ pairs.](#)

By CDF Collaboration (F. Abe et al.).

[10.1103/PhysRevLett.79.1992.](#)

Phys.Rev.Lett. 79 (1997) 1992-1997.

370) [Measurement of diffractive dijet production at the Tevatron.](#)

By CDF Collaboration (F. Abe et al.).

[10.1103/PhysRevLett.79.2636.](#)

Phys.Rev.Lett. 79 (1997) 2636-2641.

371) [Production of \$J/\psi\$ mesons from \$\chi_c\$ meson decays in \$p\bar{p}\$ collisions at \$\sqrt{s} = 1.8\$ TeV.](#)

By CDF Collaboration (F. Abe et al.).

[10.1103/PhysRevLett.79.578.](#)

Phys.Rev.Lett. 79 (1997) 578-583.

372) [Search for gluinos and squarks at the Fermilab Tevatron Collider.](#)

By CDF Collaboration (F. Abe et al.).

[10.1103/PhysRevD.56.R1357.](#)

Phys.Rev. D56 (1997) 1357-1362.

373) [\$J/\psi\$ and \$\psi\(2S\)\$ production in \$p\bar{p}\$ collisions at \$\sqrt{s} = 1.8\$ TeV.](#)

By CDF Collaboration (F. Abe et al.).

[10.1103/PhysRevLett.79.572.](#)

Phys.Rev.Lett. 79 (1997) 572-577.

374) [Observation of diffractive \$W\$ boson production at the Tevatron.](#)

By CDF Collaboration (F. Abe et al.).

[hep-ex/9703010].

[10.1103/PhysRevLett.78.2698.](#)

Phys.Rev.Lett. 78 (1997) 2698-2703.

375) [Search for new particles decaying to dijets at CDF.](#)

By CDF Collaboration (F. Abe et al.).

[hep-ex/9702004].

[10.1103/PhysRevD.55.R5263.](#)

Phys.Rev. D55 (1997) 5263-5268.

376) [ATLAS calorimeter performance Technical Design Report.](#)

By ATLAS Collaboration (A. Airapetian et al.).

377) [ATLAS computing technical proposal.](#)

By ATLAS Collaboration (A. Airapetian et al.).

378) [The Collider Detector at Fermilab: Collected physics papers.](#)

By CDF Collaboration (F. Abe et al.).

379) [The CDF-II detector: Technical design report.](#)

By CDF Collaboration (R. Blair et al.).

380) [Search for third generation leptoquarks in \$\bar{p}p\$ collisions at \$\sqrt{s} = 1.8\$ TeV.](#)

By CDF Collaboration (F. Abe et al.).

[10.1103/PhysRevLett.78.2906.](#)

Phys.Rev.Lett. 78 (1997) 2906-2911.

381) [Measurement of dijet angular distributions at CDF.](#)

By CDF Collaboration (F. Abe et al.).

[hep-ex/9609011].

[10.1103/PhysRevLett.77.5336.](#)

Phys.Rev.Lett. 77 (1996) 5336-5341, Erratum-ibid. 78 (1997) 4307.

382) [Observation of \$W^+ W^-\$ production in \$\bar{p}p\$ collisions at \$\sqrt{s} = 1.8\$ TeV.](#)

By CDF Collaboration (F. Abe et al.).

[10.1103/PhysRevLett.78.4536.](#)

Phys.Rev.Lett. 78 (1997) 4536-4540.

383) [Measurement of the branching fraction \$B\(B \rightarrow J/\psi \pi^+\)\$ and search for \$B \rightarrow J/\psi \pi^+\$.](#)

By CDF Collaboration (F. Abe et al.).

[10.1103/PhysRevLett.77.5176.](#)

Phys.Rev.Lett. 77 (1996) 5176-5181.

384) [Observation of \$\Lambda_b^0 \rightarrow J/\psi \Lambda\$ at the Fermilab proton antiproton collider.](#)

By CDF Collaboration (F. Abe et al.).

[10.1103/PhysRevD.55.1142.](#)

Phys.Rev. D55 (1997) 1142-1152.

385) [Measurement of \$b\bar{b}\$ production correlations, \$B^0 \rightarrow \bar{B}^0\$ mixing, and a limit on \$\epsilon\(B\)\$ in \$p\bar{p}\$ collisions at \$\sqrt{s} = 1.8\$ TeV.](#)

By CDF Collaboration (F. Abe et al.).

[10.1103/PhysRevD.55.2546.](#)

Phys.Rev. D55 (1997) 2546-2558.

386) [Measurement of the \$\gamma + D^{*\pm}\$ cross section in \$\bar{p}p\$ collisions at \$\sqrt{s} = 1.8\$ TeV.](#)

By CDF Collaboration (F. Abe et al.).

[10.1103/PhysRevLett.77.5005.](#)

Phys.Rev.Lett. 77 (1996) 5005-5010.

387) [Ratios of bottom meson branching fractions involving \$J/\psi\$ mesons and determination of \$b\$ quark fragmentation fractions.](#)

By CDF Collaboration (F. Abe et al.).

[hep-ex/9607003].

[10.1103/PhysRevD.54.6596.](#)

Phys.Rev. D54 (1996) 6596-6609.

388) [Further properties of high mass multi - jet events at the Fermilab proton - anti-proton collider.](#)

By CDF Collaboration (F. Abe et al.).

[hep-ex/9605004].

[10.1103/PhysRevD.54.4221.](#)

Phys.Rev. D54 (1996) 4221-4233.

389) [Measurement of the lifetime of the \$B_s^0\$ meson using the exclusive decay mode \$B_s^0 \rightarrow J/\psi \psi\$.](#)

By CDF Collaboration (F. Abe et al.).

[10.1103/PhysRevLett.77.1945.](#)

Phys.Rev.Lett. 77 (1996) 1945-1949.

390) [Measurement of \$\Lambda_b^0\$ lifetime using \$\Lambda_b^0 \rightarrow \Lambda_c^+ \ell \bar{\nu}_\ell\$.](#)

By CDF Collaboration (F. Abe et al.).

[10.1103/PhysRevLett.77.1439.](#)

Phys.Rev.Lett. 77 (1996) 1439-1443.

391) [Forward-backward charge asymmetry of electron pairs above the \$Z^0\$ pole.](#)

By CDF Collaboration (F. Abe et al.).

[10.1103/PhysRevLett.77.2616.](#)

Phys.Rev.Lett. 77 (1996) 2616-2621.

392) [Properties of jets in Z boson events from 1.8 TeV \$\bar{p}p\$ collisions.](#)

By CDF Collaboration (F. Abe et al.).

[hep-ex/9603003].

[10.1103/PhysRevLett.77.448.](#)

Phys.Rev.Lett. 77 (1996) 448-453.

393) [Search for flavor changing neutral current \$B\$ meson decays in \$\bar{p}p\$ collisions at \$\sqrt{s} = 1.8\$ TeV.](#)

By CDF Collaboration (F. Abe et al.).

[10.1103/PhysRevLett.76.4675.](#)

Phys.Rev.Lett. 76 (1996) 4675-4680.

394) [Measurement of the \$B^+\$ and \$\bar{B}^0\$ meson lifetimes using semileptonic decays.](#)

By CDF Collaboration (F. Abe et al.).

[10.1103/PhysRevLett.76.4462.](#)

Phys.Rev.Lett. 76 (1996) 4462-4467.

395) [Search for chargino - neutralino production in \$\bar{p}p\$ collisions at \$\sqrt{s} = 1.8\$ TeV.](#)

By CDF Collaboration (F. Abe et al.).

[hep-ex/9603001].

[10.1103/PhysRevLett.76.4307.](#)

Phys.Rev.Lett. 76 (1996) 4307-4311.

396) [Inclusive jet cross section in \$\bar{p}p\$ collisions at \$\sqrt{s}=1.8\$ TeV.](#)

By CDF Collaboration (F. Abe et al.).

[hep-ex/9601008].

[10.1103/PhysRevLett.77.438.](#)

Phys.Rev.Lett. 77 (1996) 438-443.

397) [Status report on CLUE.](#)

By D. Alexandreas, B. Bartoli, D. Bastieri, Franco Bedeschi, E. Bertolucci, C. Bigongiari, R. Biral, G. Busetto et al..

[10.1016/0168-9002\(94\)01622-4](https://doi.org/10.1016/0168-9002(94)01622-4).

Nucl.Instrum.Meth. A360 (1995) 385-389.

398) [Search for the rare decay \$W^{\pm} \rightarrow \pi^{\pm} + \gamma\$.](#)

By CDF Collaboration (F. Abe et al.).

[10.1103/PhysRevLett.76.2852](https://doi.org/10.1103/PhysRevLett.76.2852).

Phys.Rev.Lett. 76 (1996) 2852-2857.

399) [Search for charged Higgs decays of the top quark using hadronic tau decays.](#)

By CDF Collaboration (F. Abe et al.).

[hep-ex/9601003].

[10.1103/PhysRevD.54.735](https://doi.org/10.1103/PhysRevD.54.735).

Phys.Rev. D54 (1996) 735-742.

400) [Search for gluino and squark cascade decays at the Fermilab Tevatron Collider.](#)

By CDF Collaboration (F. Abe et al.).

[10.1103/PhysRevLett.76.2006](https://doi.org/10.1103/PhysRevLett.76.2006).

Phys.Rev.Lett. 76 (1996) 2006-2010.

401) [Reconstruction of \$B^0 \rightarrow J/\psi K_s^0\$ and measurement of ratios of branching ratios involving \$B \rightarrow J/\psi K^*\$.](#)

By CDF Collaboration (F. Abe et al.).

[10.1103/PhysRevLett.76.2015](https://doi.org/10.1103/PhysRevLett.76.2015).

Phys.Rev.Lett. 76 (1996) 2015-2020.

402) [Measurement of the mass of the \$B_s^0\$ meson.](#)

By CDF Collaboration (F. Abe et al.).

[10.1103/PhysRevD.53.3496](https://doi.org/10.1103/PhysRevD.53.3496).

Phys.Rev. D53 (1996) 3496-3505.

403) [Measurement of \$\sigma \cdot B\(W \rightarrow e \nu\)\$ and \$\sigma \cdot B\(Z^0 \rightarrow e^+ e^-\)\$ in \$p\bar{p}\$ collisions at \$\sqrt{s} = 1.8\$ TeV.](#)

By CDF Collaboration (F. Abe et al.).

[hep-ex/9509010].

[10.1103/PhysRevLett.76.3070](https://doi.org/10.1103/PhysRevLett.76.3070).

Phys.Rev.Lett. 76 (1996) 3070-3075.

404) [Measurement of correlated \$\mu - \bar{b}\$ jet cross-sections in \$p\bar{p}\$ collisions at \$\sqrt{s} = 1.8\$ TeV.](#)

By CDF Collaboration (F. Abe et al.).

[hep-ex/9508017].

[10.1103/PhysRevD.53.1051](https://doi.org/10.1103/PhysRevD.53.1051).

Phys.Rev. D53 (1996) 1051-1065.

405) [\$\Upsilon\$ production in \$p\bar{p}\$ collisions at \$\sqrt{s} = 1.8\$ TeV.](#)

By CDF Collaboration (F. Abe et al.).

[10.1103/PhysRevLett.75.4358](https://doi.org/10.1103/PhysRevLett.75.4358).

Phys.Rev.Lett. 75 (1995) 4358.

406) [Measurement of the polarization in the decays \$B_d \rightarrow J/\psi K^{*0}\$ and \$B_s \rightarrow J/\psi \phi\$.](#)

By CDF Collaboration (F. Abe et al.).

[10.1103/PhysRevLett.75.3068](https://doi.org/10.1103/PhysRevLett.75.3068).

Phys.Rev.Lett. 75 (1995) 3068.

407) [Study of \$t\bar{t}\$ production \$p\bar{p}\$ collisions using total transverse energy.](#)

By CDF Collaboration (F. Abe et al.).

[hep-ex/9506006].

[10.1103/PhysRevLett.75.3997.](#)

Phys.Rev.Lett. 75 (1995) 3997.

408) [A Measurement of the ratio \$\sigma \times B\(p\bar{p} \rightarrow W \rightarrow e \nu\) / \sigma \times B\(p\bar{p} \rightarrow Z^0 \rightarrow e e\)\$ in \$p\bar{p}\$ collisions at \$\sqrt{s} = 1800\$ GeV.](#)

By CDF Collaboration (F. Abe et al.).

[10.1103/PhysRevD.52.2624.](#)

Phys.Rev. D52 (1995) 2624-2655.

409) [Identification of top quarks at CDF using kinematic variables.](#)

By CDF Collaboration (F. Abe et al.).

[10.1103/PhysRevD.52.R2605.](#)

Phys.Rev. D52 (1995) 2605-2609.

410) [A Search for second generation leptoquarks in \$p\bar{p}\$ collisions at \$\sqrt{s} = 1.8\$ TeV.](#)

By CDF Collaboration (F. Abe et al.).

[10.1103/PhysRevLett.75.1012.](#)

Phys.Rev.Lett. 75 (1995) 1012-1016.

411) [Measurement of the \$B\$ meson differential cross-section, \$d\sigma / dp_T\$, in \$p\bar{p}\$ collisions at \$\sqrt{s} = 1.8\$ TeV.](#)

By CDF Collaboration (F. Abe et al.).

[hep-ex/9503013].

[10.1103/PhysRevLett.75.1451.](#)

Phys.Rev.Lett. 75 (1995) 1451-1455.

412) [Properties of high mass multi - jet events at the Fermilab \$p\bar{p}\$ collider.](#)

By CDF Collaboration (F. Abe et al.).

[10.1103/PhysRevLett.75.608.](#)

Phys.Rev.Lett. 75 (1995) 608-612.

413) [Measurement of the \$W\$ boson mass.](#)

By CDF Collaboration (F. Abe et al.).

[hep-ex/9503007].

[10.1103/PhysRevLett.75.11.](#)

Phys.Rev.Lett. 75 (1995) 11-16.

414) [Measurement of the \$W\$ boson mass.](#)

By CDF Collaboration (F. Abe et al.).

[10.1103/PhysRevD.52.4784.](#)

Phys.Rev. D52 (1995) 4784-4827.

415) [Limits on \$W W Z\$ and \$W W \gamma\$ couplings from \$W W\$ and \$W Z\$ production in \$p\bar{p}\$ collisions at \$\sqrt{s} = 1.8\$ TeV.](#)

By CDF Collaboration (F. Abe et al.).

[hep-ex/9503009].

[10.1103/PhysRevLett.75.1017.](#)

Phys.Rev.Lett. 75 (1995) 1017-1022.

416) [Observation of top quark production in \$p\bar{p}\$ collisions.](#)

By CDF Collaboration (F. Abe et al.).

[hep-ex/9503002].

[10.1103/PhysRevLett.74.2626.](#)

Phys.Rev.Lett. 74 (1995) 2626-2631.

417) [Measurement of the \$B_s\$ meson lifetime.](#)

By CDF Collaboration (F. Abe et al.).

[hep-ex/9412017].

[10.1103/PhysRevLett.74.4988.](#)

Phys.Rev.Lett. 74 (1995) 4988.

418) [Search for new particles decaying to dijets in \$p\bar{p}\$ collisions at \$\sqrt{s} = 1.8\$ TeV.](#)

By CDF Collaboration (F. Abe et al.).

[hep-ex/9501001].

[10.1103/PhysRevLett.74.3538.](#)

Phys.Rev.Lett. 74 (1995) 3538-3543.

419) [Measurement of the total elastic and single diffractive cross-sections at \$s^{1/2} = 546\$ -GeV and 1800-GeV using the luminosity independent method.](#)

By CDF Collaboration (R. Paoletti for the collaboration).

In *Meribel les Allues 1994, QCD and high energy hadronic interactions* 303-308.

420) [Search for the top quark decaying to a charged Higgs boson in \$\bar{p}p\$ collisions at \$\sqrt{s} = 1.8\$ TeV.](#)

By CDF Collaboration (F. Abe et al.).

[10.1103/PhysRevLett.73.2667.](#)

Phys.Rev.Lett. 73 (1994) 2667-2671.

421) [Measurement of the \$B\$ meson and \$b\$ quark cross-sections at \$\sqrt{s} = 1.8\$ TeV using the exclusive decay \$B^0 \rightarrow J/\psi K^*0\$ \(892\).](#)

By CDF Collaboration (F. Abe et al.).

[10.1103/PhysRevD.50.4252.](#)

Phys.Rev. D50 (1994) 4252.

422) [Search for charged bosons heavier than the \$W\$ in \$p\bar{p}\$ collisions at \$\sqrt{s} = 1800\$ GeV.](#)

By CDF Collaboration (F. Abe et al.).

[10.1103/PhysRevLett.74.2900.](#)

Phys.Rev.Lett. 74 (1995) 2900-2904.

423) [Kinematic evidence for top quark pair production in \$W\$ + multi - jet events in \$p\bar{p}\$ collisions at \$\sqrt{s} = 1.8\$ TeV.](#)

By CDF Collaboration (F. Abe et al.).

[hep-ex/9412009].

[10.1103/PhysRevD.51.4623.](#)

Phys.Rev. D51 (1995) 4623-4637.

424) [\$b\bar{b}\$ quark pair correlations in \$p\bar{p}\$ collisions at \$S^{1/2} = 1.8\$ -TeV.](#)

By CDF Collaboration (F. Abe et al.).

Unpublished - incorporated into Phys.Rev. D55, 2546 (1997).

425) [Limits on \$Z\$ -photon couplings from \$p\bar{p}\$ interactions at \$\sqrt{s} = 1.8\$ TeV.](#)

By CDF Collaboration (F. Abe et al.).

[10.1103/PhysRevLett.74.1941.](#)

Phys.Rev.Lett. 74 (1995) 1941-1945.

426) [The Charge asymmetry in \$W\$ boson decays produced in \$p\bar{p}\$ collisions at \$\sqrt{s} = 1.8\$ TeV.](#)

By CDF Collaboration (F. Abe et al.).

[hep-ex/9501008].

[10.1103/PhysRevLett.74.850.](#)

Phys.Rev.Lett. 74 (1995) 850-854.

427) [A Direct measurement of the \$W\$ boson width.](#)

By CDF Collaboration (F. Abe et al.).

[10.1103/PhysRevLett.74.341.](#)

Phys.Rev.Lett. 74 (1995) 341-345.

428) [Measurement of \$W - \gamma\$ couplings with {CDF} in \$p\bar{p}\$ collisions at \$\sqrt{s} = 1.8\$ TeV.](#)

By CDF Collaboration (F. Abe et al.).

[10.1103/PhysRevLett.74.1936.](#)

Phys.Rev.Lett. 74 (1995) 1936-1940.

429) [Measurement of \$W + \gamma\$ and \$Z + \gamma\$ cross-sections in the electron and muon channels in \$S^{\(1/2\)} = 1.8\$ -TeV anti-P P collisions.](#)

By CDF Collaboration (F. Abe et al.).

430) [Search for radiative decays of neutralinos in \$p\bar{p}\$ collisions at \$\sqrt{s} = 1.8\$ TeV.](#)

By CDF Collaboration (F. Abe et al.).

[10.1103/PhysRevLett.75.613.](#)

Phys.Rev.Lett. 75 (1995) 613-617.

431) [Observation of rapidity gaps in \$\bar{p}p\$ collisions at 1.8 TeV.](#)

By CDF Collaboration (F. Abe et al.).

[10.1103/PhysRevLett.74.855.](#)

Phys.Rev.Lett. 74 (1995) 855-859.

432) [A Precision measurement of the prompt photon cross-section in \$p\bar{p}\$ collisions at \$\sqrt{s} = 1.8\$ TeV.](#)

By CDF Collaboration (F. Abe et al.).

[10.1103/PhysRevLett.73.2662.](#)

Phys.Rev.Lett. 73 (1994) 2662-2666, Erratum-ibid. 74 (1995) 1891-1893.

433) [Search for new gauge bosons decaying into dielectrons in \$\bar{p}p\$ collisions at \$\sqrt{s}=1.8\$ TeV.](#)

By CDF Collaboration (F. Abe et al.).

[10.1103/PhysRevD.51.R949.](#)

Phys.Rev. D51 (1995) 949-953.

434) [Analysis of jet charged particle momentum distributions for quark - gluon separation in \$\bar{p} p\$ collisions at \$\sqrt{s} = 1.8\$ -TeV.](#)

By CDF Collaboration (F. Abe et al.).

435) [Evidence for color coherence in \$p\bar{p}\$ collisions at \$\sqrt{s} = 1.8\$ TeV.](#)

By CDF Collaboration (F. Abe et al.).

[10.1103/PhysRevD.50.5562.](#)

Phys.Rev. D50 (1994) 5562-5579.

436) [\$W\$ boson + jet angular distribution in \$p\bar{p}\$ collisions at \$\sqrt{s} = 1.8\$ TeV.](#)

By CDF Collaboration (F. Abe et al.).

[10.1103/PhysRevLett.73.2296.](#)

Phys.Rev.Lett. 73 (1994) 2296-2300.

437) [Measurement of the ratio \$\sigma_B\(W \rightarrow e \nu\) / \sigma_B\(Z^0 \rightarrow e^+ e^-\)\$ in \$\bar{p}p\$](#)

[collisions at \$\sqrt{s} = 1.8\$ TeV.](#)

By CDF Collaboration (F. Abe et al.).

[10.1103/PhysRevLett.73.220.](#)

Phys.Rev.Lett. 73 (1994) 220-224.

438) [Evidence for top quark production in \$\bar{p}p\$ collisions at \$\sqrt{s} = 1.8\$ TeV.](#)

By CDF Collaboration (F. Abe et al.).

[hep-ex/9405005].

[10.1103/PhysRevLett.73.225.](#)

Phys.Rev.Lett. 73 (1994) 225-231.

439) [Evidence for top quark production in \$\bar{p}p\$ collisions at \$\sqrt{s} = 1.8\$ TeV.](#)

By CDF Collaboration (F. Abe et al.).

[10.1103/PhysRevD.50.2966.](#)

Phys.Rev. D50 (1994) 2966-3026.

440) [Measurement of the \$B^+\$ and \$B^0\$ meson lifetimes.](#)

By CDF Collaboration (F. Abe et al.).

[10.1103/PhysRevLett.72.3456.](#)

Phys.Rev.Lett. 72 (1994) 3456-3460.

441) [The CLUE mirrors.](#)

By D. Alexandreas, B. Bartoli, D. Bastieri, Franco Bedeschi, E. Bertolucci, P. Bicchi, G. Chiarelli, R. Conte et al..

In *Calgary 1993, Towards a major atmospheric Cherenkov detector for TeV astro/particle physics* 169-175.

442) [Status report on CLUE.](#)

By D. Alexandreas, B. Bartoli, Franco Bedeschi, E. Bertolucci, P. Bicchi, C. Bigongiari, G. Busetto, S. Centro et al..

In *Calgary 1993, Towards a major atmospheric Cherenkov detector for TeV astro/particle physics* 49-55.

443) [A Search for the top quark decaying to charged Higgs in \$p\bar{p}\$ collisions at \$\sqrt{s} = 1.8\$ TeV.](#)

By CDF Collaboration (F. Abe et al.).

[10.1103/PhysRevLett.72.1977.](#)

Phys.Rev.Lett. 72 (1994) 1977-1981.

444) [Search for excited quarks in \$p\bar{p}\$ collisions at \$\sqrt{s} = 1.8\$ TeV.](#)

By CDF Collaboration (F. Abe et al.).

[10.1103/PhysRevLett.72.3004.](#)

Phys.Rev.Lett. 72 (1994) 3004-3008.

445) [Measurement of the \$\bar{p}p\$ total cross-section at \$\sqrt{s} = 546\$ GeV and 1800-GeV.](#)

By CDF Collaboration (F. Abe et al.).

[10.1103/PhysRevD.50.5550.](#)

Phys.Rev. D50 (1994) 5550-5561.

446) [Measurement of small angle \$\bar{p}p\$ elastic scattering at \$\sqrt{s} = 546\$ GeV and 1800 GeV.](#)

By CDF Collaboration (F. Abe et al.).

[10.1103/PhysRevD.50.5518.](#)

Phys.Rev. D50 (1994) 5518-5534.

447) [Measurement of \$\bar{p}p\$ single diffraction dissociation at \$\sqrt{s} = 546\$ GeV and 1800 GeV.](#)

By CDF Collaboration (F. Abe et al.).

[10.1103/PhysRevD.50.5535](https://arxiv.org/abs/10.1103/PhysRevD.50.5535).

Phys.Rev. D50 (1994) 5535-5549.

448) [Top dilepton search at CDF.](#)

By CDF Collaboration (F. Abe et al.).

449) [Probing the gluon distribution with the SS - OS dijet cross-section ratio.](#)

By CDF Collaboration (F. Abe et al.).

450) [Measurement of inclusive jet cross-sections in \$p\bar{p}\$ collisions at CDF.](#)

By CDF Collaboration (F. Abe et al.).

451) [SUSY search using trilepton events from \$p\bar{p}\$ collisions at \$\sqrt{s}=1.8\$ -TeV.](#)

By CDF Collaboration (F. Abe et al.).

452) [Measurement of the \$B^+\$ and \$B^0\$ lifetimes.](#)

By CDF Collaboration (F. Abe et al.).

453) [The Lepton charge asymmetry in the decay of \$W\$ bosons produced in \$p\bar{p}\$ collisions at \$\sqrt{s}=1.8\$ -TeV.](#)

By CDF Collaboration (F. Abe et al.).

454) [Measurement of the B meson and b quark cross-sections in \$p\bar{p}\$ collisions at \$\sqrt{s}=1.8\$ -TeV using the exclusive decays \$B^+ \rightarrow J/\psi K^+\$ and \$B^0 \rightarrow J/\psi K^0\$.](#)

By CDF Collaboration (F. Abe et al.).

455) [The Two jet differential cross-section at CDF.](#)

By CDF Collaboration (F. Abe et al.).

456) [Search for excited quarks in \$p\bar{p}\$ collisions at \$\sqrt{s}=1.8\$ -TeV.](#)

By CDF Collaboration (F. Abe et al.).

457) [The Cross-section for the production of \$b\bar{b}\$ pairs in \$p\bar{p}\$ collisions at \$\sqrt{s}=1.8\$ -TeV.](#)

By CDF Collaboration (F. Abe et al.).

458) [A Study of events with the highest total transverse energy in CDF.](#)

By CDF Collaboration (F. Abe et al.).

459) [Direct photon results from CDF.](#)

By CDF Collaboration (F. Abe et al.).

460) [The Center-of-mass angular distribution of prompt photons produced in \$p\$ anti- \$p\$ collisions at \$\sqrt{s}=1.8\$ -TeV.](#)

By CDF Collaboration (F. Abe et al.).

461) [A Search for new gauge bosons in \$p\bar{p}\$ collisions at \$\sqrt{s}=1.8\$ -TeV in the dielectron decay mode.](#)

By CDF Collaboration (F. Abe et al.).

462) [Study of four jet events and evidence for double parton interactions in \$p\bar{p}\$ collisions at \$\sqrt{s}=1.8\$ TeV.](#)

By CDF Collaboration (F. Abe et al.).

[10.1103/PhysRevD.47.4857](https://arxiv.org/abs/10.1103/PhysRevD.47.4857).

Phys.Rev. D47 (1993) 4857-4871.

- 463) [Inclusive \$\chi_c\$ and \$b\$ quark production in \$\bar{p}p\$ collisions at \$\sqrt{s} = 1.8\$ TeV.](#)
By CDF Collaboration (F. Abe et al.).
[10.1103/PhysRevLett.71.2537.](#)
Phys.Rev.Lett. 71 (1993) 2537-2541.
- 464) [Search for quark compositeness, axiguons and heavy particles using the dijet invariant mass spectrum observed in \$\bar{p}p\$ collisions.](#)
By CDF Collaboration (F. Abe et al.).
[10.1103/PhysRevLett.71.2542.](#)
Phys.Rev.Lett. 71 (1993) 2542-2546.
- 465) [Measurement of the average lifetime of B hadrons produced in \$\bar{p}p\$ collisions at \$\sqrt{s} = 1.8\$ TeV.](#)
By CDF Collaboration (F. Abe et al.).
[10.1103/PhysRevLett.71.3421.](#)
Phys.Rev.Lett. 71 (1993) 3421-3426.
- 466) [Measurement of bottom quark production in 1.8 TeV \$\bar{p}p\$ collisions using semileptonic decay muons.](#)
By CDF Collaboration (F. Abe et al.).
[10.1103/PhysRevLett.71.2396.](#)
Phys.Rev.Lett. 71 (1993) 2396-2400.
- 467) [Measurement of Drell-Yan electron and muon pair differential cross-sections in \$\bar{p}p\$ collisions at \$\sqrt{s} = 1.8\$ TeV.](#)
By CDF Collaboration (F. Abe et al.).
[10.1103/PhysRevD.49.R1.](#)
Phys.Rev. D49 (1994) 1-6.
- 468) [Observation of the decay \$B_s^0 \rightarrow J/\psi \phi\$ in \$\bar{p}p\$ collisions at \$\sqrt{s} = 1.8\$ TeV.](#)
By CDF Collaboration (F. Abe et al.).
[10.1103/PhysRevLett.71.1685.](#)
Phys.Rev.Lett. 71 (1993) 1685-1689.
- 469) [A Search for first generation leptoquarks in \$\bar{p}p\$ collisions at \$\sqrt{s} = 1.8\$ TeV.](#)
By CDF Collaboration (F. Abe et al.).
[10.1103/PhysRevD.48.R3939.](#)
Phys.Rev. D48 (1993) 3939-3944.
- 470) [Measurement of the bottom quark production cross-section using semileptonic decay electrons in \$\bar{p}p\$ collisions at \$\sqrt{s} = 1.8\$ TeV.](#)
By CDF Collaboration (F. Abe et al.).
[10.1103/PhysRevLett.71.500.](#)
Phys.Rev.Lett. 71 (1993) 500-504.
- 471) [Measurement of jet multiplicity in W events produced in \$\bar{p}p\$ collisions at \$\sqrt{s} = 1.8\$ -
 \$1.8\$ TeV.](#)
By CDF Collaboration (F. Abe et al.).
[10.1103/PhysRevLett.70.4042.](#)
Phys.Rev.Lett. 70 (1993) 4042-4046.
- 472) [The Center-of-mass angular distribution of prompt photons produced in \$\bar{p}p\$ collisions at \$\sqrt{s} = 1.8\$ TeV.](#)
By CDF Collaboration (F. Abe et al.).
[10.1103/PhysRevLett.71.679.](#)
Phys.Rev.Lett. 71 (1993) 679-683.

- 473) [Measurement of the dijet mass distribution in \$p\bar{p}\$ collisions at \$\sqrt{s} = 1.8\$ TeV.](#)
By CDF Collaboration (F. Abe et al.).
[10.1103/PhysRevD.48.998.](#)
Phys.Rev. D48 (1993) 998-1008.
- 474) [A Prompt photon cross-section measurement in \$\bar{p}p\$ collisions at \$\sqrt{s} = 1.8\$ TeV.](#)
By CDF Collaboration (F. Abe et al.).
[10.1103/PhysRevD.48.2998.](#)
Phys.Rev. D48 (1993) 2998-3025.
- 475) [A Background - free detector for cosmic ray showers in the atmosphere.](#)
By A. Menzione, F. Angelini, Franco Bedeschi, R. Bellazzini, E. Bertolucci, G. Chiarelli, M. Mariotti, M.M. Massai et al..
[10.1016/0168-9002\(92\)90714-F.](#)
Nucl.Instrum.Meth. A315 (1992) 267-271.
- 476) [Measurement of the cross-section for production of two isolated prompt photons in \$\bar{p}p\$ collisions at \$\sqrt{s} = 1.8\$ TeV.](#)
By CDF Collaboration (F. Abe et al.).
[10.1103/PhysRevLett.70.2232.](#)
Phys.Rev.Lett. 70 (1993) 2232-2236.
- 477) [On the surjectivity of Wahl maps on general curve.](#)
By Roberto Paoletti.
[alg-geom/9302002].
- 478) [Comparison of jet production in \$\bar{p}p\$ collisions at \$\sqrt{s} = 546\$ GeV and 1800 GeV.](#)
By CDF Collaboration (F. Abe et al.).
[10.1103/PhysRevLett.70.1376.](#)
Phys.Rev.Lett. 70 (1993) 1376-1380.
- 479) [Search for \$\Lambda_b \rightarrow J/\psi \Lambda^0\$ in \$p\bar{p}\$ collisions at \$\sqrt{s} = 1.8\$ TeV.](#)
By CDF Collaboration (F. Abe et al.).
[10.1103/PhysRevD.47.R2639.](#)
Phys.Rev. D47 (1993) 2639-2643.
- 480) [Inclusive \$J/\psi\$, \$\psi\(2S\)\$ and \$b\bar{b}\$ quark production in \$\bar{p}p\$ collisions at \$\sqrt{s} = 1.8\$ TeV.](#)
By CDF Collaboration (F. Abe et al.).
[10.1103/PhysRevLett.69.3704.](#)
Phys.Rev.Lett. 69 (1992) 3704-3708.
- 481) [A Measurement of jet shapes in \$p\bar{p}\$ collisions at \$\sqrt{s} = 1.8\$ TeV.](#)
By CDF Collaboration (F. Abe et al.).
[10.1103/PhysRevLett.70.713.](#)
Phys.Rev.Lett. 70 (1993) 713-717.
- 482) [Search for squarks and gluinos from \$\bar{p}p\$ collisions at \$\sqrt{s} = 1.8\$ TeV.](#)
By CDF Collaboration (F. Abe et al.).
[10.1103/PhysRevLett.69.3439.](#)
Phys.Rev.Lett. 69 (1992) 3439-3443.
- 483) [Limit on the rare decay \$W^+ \rightarrow \gamma \pi^+ \pi^0\$ in \$p\bar{p}\$ collisions at \$\sqrt{s} = 1.8\$ TeV.](#)
By CDF Collaboration (F. Abe et al.).

[10.1103/PhysRevLett.69.2160.](#)

Phys.Rev.Lett. 69 (1992) 2160-2163.

484) [The Dijet angular distribution in \$p\bar{p}\$ collisions at \$\sqrt{s} = 1.8\$ TeV.](#)

By CDF Collaboration (F. Abe et al.).

[10.1103/PhysRevLett.69.2896.](#)

Phys.Rev.Lett. 69 (1992) 2896-2900.

485) [Limits on the production of massive stable charged particles.](#)

By CDF Collaboration (F. Abe et al.).

[10.1103/PhysRevD.46.R1889.](#)

Phys.Rev. D46 (1992) 1889-1894.

486) [A Measurement of the production and muonic decay rate of \$W\$ and \$Z\$ bosons in \$p\bar{p}\$ collisions at \$\sqrt{s} = 1.8\$ TeV.](#)

By CDF Collaboration (F. Abe et al.).

[10.1103/PhysRevLett.69.28.](#)

Phys.Rev.Lett. 69 (1992) 28-32.

487) [Measurement of the isolated prompt photon cross-sections in \$p\bar{p}\$ collisions at \$\sqrt{s} = 1.8\$ TeV.](#)

By CDF Collaboration (F. Abe et al.).

[10.1103/PhysRevLett.68.2734.](#)

Phys.Rev.Lett. 68 (1992) 2734-2738.

488) [A Measurement of the \$B\$ meson and \$b\$ quark cross-sections at \$\sqrt{s} = 1.8\$ TeV using the exclusive decay \$B^{\pm} \rightarrow J/\psi K^{\pm}\$.](#)

By CDF Collaboration (F. Abe et al.).

[10.1103/PhysRevLett.68.3403.](#)

Phys.Rev.Lett. 68 (1992) 3403-3407.

489) [Measurement of \$B^0\$ mixing at the Fermilab Tevatron collider.](#)

By CDF Collaboration (F. Abe et al.).

[10.1103/PhysRevLett.67.3351.](#)

Phys.Rev.Lett. 67 (1991) 3351-3355.

490) [Measurement of the ratio \$\sigma_B\(W \rightarrow \nu \tau\) / \sigma_B\(W \rightarrow e \nu\)\$, in \$p\bar{p}\$ collisions at \$\sqrt{s} = 1.8\$ TeV.](#)

By CDF Collaboration (F. Abe et al.).

[10.1103/PhysRevLett.68.3398.](#)

Phys.Rev.Lett. 68 (1992) 3398-3402.

491) [A Limit on the top quark mass from \$p\bar{p}\$ collisions at \$\sqrt{s} = 1.8\$ TeV.](#)

By CDF Collaboration (F. Abe et al.).

[10.1103/PhysRevD.45.3921.](#)

Phys.Rev. D45 (1992) 3921-3948.

492) [Lepton asymmetry in \$W\$ decays from \$p\bar{p}\$ collisions at \$\sqrt{s} = 1.8\$ TeV.](#)

By CDF Collaboration (F. Abe et al.).

[10.1103/PhysRevLett.68.1458.](#)

Phys.Rev.Lett. 68 (1992) 1458-1462.

493) [Measurement of the \$Z \rightarrow T\$ distribution in \$p\bar{p}\$ collisions at \$\sqrt{s} = 1.8\$ TeV.](#)

By CDF Collaboration (F. Abe et al.).

[10.1103/PhysRevLett.67.2937.](#)

Phys.Rev.Lett. 67 (1991) 2937-2941.

- 494) [Properties of events with large total transverse energy produced in \$p\bar{p}\$ collisions at \$\sqrt{s} = 1.8\$ TeV.](#)
By CDF Collaboration (F. Abe et al.).
[10.1103/PhysRevD.45.2249.](#)
Phys.Rev. D45 (1992) 2249-2263.
- 495) [A Search for new gauge bosons in \$p\bar{p}\$ collisions at \$\sqrt{s} = 1.8\$ TeV.](#)
By CDF Collaboration (F. Abe et al.).
[10.1103/PhysRevLett.68.1463.](#)
Phys.Rev.Lett. 68 (1992) 1463-1467.
- 496) [A Lower limit on the top quark mass from events with two leptons in \$p\bar{p}\$ collisions at \$\sqrt{s} = 1.8\$ TeV.](#)
By CDF Collaboration (F. Abe et al.).
[10.1103/PhysRevLett.68.447.](#)
Phys.Rev.Lett. 68 (1992) 447-451.
- 497) [SVX test beam results.](#)
By Franco Bedeschi, S. Dell'Agello, M. Mariotti, A. Menzione, R. Paoletti, F. Raffaelli, Giuseppe Francesco Tartarelli, N. Turini et al..
[10.1016/0168-9002\(92\)90702-6.](#)
Nucl.Instrum.Meth. A315 (1992) 188-192.
- 498) [The Topology of three jet events in \$p\bar{p}\$ collisions at \$\sqrt{s} = 1.8\$ TeV.](#)
By CDF Collaboration (F. Abe et al.).
[10.1103/PhysRevD.45.1448.](#)
Phys.Rev. D45 (1992) 1448-1458.
- 499) [Inclusive jet cross-section in \$p\bar{p}\$ collisions at \$\sqrt{s} = 1.8\$ TeV.](#)
By CDF Collaboration (F. Abe et al.).
[10.1103/PhysRevLett.68.1104.](#)
Phys.Rev.Lett. 68 (1992) 1104-1108.
- 500) [Search for \$W^{\prime} \rightarrow e \nu\$ and \$W^{\prime} \rightarrow \mu \nu\$ in \$p\bar{p}\$ collisions at \$\sqrt{s} = 1.8\$ TeV.](#)
By CDF Collaboration (F. Abe et al.).
[10.1103/PhysRevLett.67.2609.](#)
Phys.Rev.Lett. 67 (1991) 2609-2613.
- 501) [Measurement of the \$e^+e^-\$ invariant mass distribution in \$p\bar{p}\$ collisions at \$\sqrt{s} = 1.8\$ TeV.](#)
By CDF Collaboration (F. Abe et al.).
[10.1103/PhysRevLett.67.2418.](#)
Phys.Rev.Lett. 67 (1991) 2418-2422.
- 502) [A Determination of \$\sin^2\theta\$ from the forward - backward asymmetry in \$p\bar{p} \rightarrow Z^0 X \rightarrow e^+e^- X\$ interactions at \$\sqrt{s} = 1.8\$ TeV.](#)
By CDF Collaboration (F. Abe et al.).
[10.1103/PhysRevLett.67.1502.](#)
Phys.Rev.Lett. 67 (1991) 1502-1506.
- 503) [Measurement of QCD jet broadening in \$p\bar{p}\$ collisions at \$\sqrt{s} = 1.8\$ TeV.](#)
By CDF Collaboration (F. Abe et al.).
[10.1103/PhysRevD.44.601.](#)
Phys.Rev. D44 (1991) 601-616.

504) [Measurement of the \$W\$ \$P\(T\)\$ distribution in \$\bar{p}p\$ collisions at \$\sqrt{s} = 1.8\$ TeV.](#)

By CDF Collaboration (F. Abe et al.).

[10.1103/PhysRevLett.66.2951.](#)

Phys.Rev.Lett. 66 (1991) 2951-2955.

505) [\$B\$ physics at CDF.](#)

By CDF Collaboration (Andrew R. Baden et al.).

506) [The Small angle spectrometer of CDF.](#)

By G. Chiarelli, S. Bertolucci, M. Cordelli, P. Giromini, S. Miscetti, A. Sansoni, T.J. Chapin, N. Giokaris et al..

[10.1016/0168-9002\(90\)91506-7.](#)

Nucl.Instrum.Meth. A289 (1990) 375-383.

507) [A Measurement of \$\sigma_B\(W \rightarrow e \nu\)\$ and \$\sigma_B\(Z^0 \rightarrow e^+ e^-\)\$ in \$\bar{p}p\$ collisions at \$\sqrt{s} = 1800\$ GeV.](#)

By CDF Collaboration (F. Abe et al.).

[10.1103/PhysRevD.44.29.](#)

Phys.Rev. D44 (1991) 29-52.

508) [A Measurement of the \$W\$ boson mass in 1.8 TeV \$\bar{p}p\$ collisions.](#)

By CDF Collaboration (F. Abe et al.).

[10.1103/PhysRevD.43.2070.](#)

Phys.Rev. D43 (1991) 2070-2093.

509) [A Measurement of the \$W\$ boson mass.](#)

By CDF Collaboration (F. Abe et al.).

[10.1103/PhysRevLett.65.2243.](#)

Phys.Rev.Lett. 65 (1990) 2243-2246.

510) [Top quark search in the electron + jet channel in \$p\bar{p}\$ collisions at \$\sqrt{s} = 1.8\$ TeV.](#)

By CDF Collaboration (F. Abe et al.).

[10.1103/PhysRevD.43.664.](#)

Phys.Rev. D43 (1991) 664-686.

511) [Jet fragmentation properties of \$\bar{p}p\$ collisions at \$\sqrt{s} = 1.8\$ TeV.](#)

By CDF Collaboration (F. Abe et al.).

[10.1103/PhysRevLett.65.968.](#)

Phys.Rev.Lett. 65 (1990) 968-971.

512) [The Dijet invariant mass at the Tevatron collider.](#)

By CDF Collaboration (Mauro Dell'Orso et al.).

513) [Search For New Heavy Quarks In Electron - Muon Events At The Fermilab Tevatron Collider.](#)

By CDF Collaboration (F. Abe et al.).

[10.1103/PhysRevLett.64.147.](#)

Phys.Rev.Lett. 64 (1990) 147-151.

514) [Status of the top quark search at CDF.](#)

By CDF Collaboration (G. William Foster et al.).

AIP Conf.Proc. 196 (1989) 354-364.

515) [Performance of surface barrier silicon detectors with x,y readout.](#)

By G. Apollinari, S. Belforte, F. Bosi, E. Focardi, R. Paoletti, G. Tonelli, F. Zetti.

[10.1016/0168-9002\(89\)91112-1.](#)

Nucl.Instrum.Meth. A279 (1989) 396-401.

516) [Two-dimensional Tracking With Surface Barrier Silicon Detectors.](#)
By G. Apollinari, S. Belforte, E. Focardi, R. Paoletti, G. Tonelli, F. Zetti.
[10.1109/23.34399.](#)

IEEE Trans.Nucl.Sci. 36 (1989) 46-53.

517) [The Two jet invariant mass distribution at \$\sqrt{s} = 1.8\$ TeV.](#)

By CDF Collaboration (F. Abe et al.).

[10.1103/PhysRevD.41.1722.](#)

Phys.Rev. D41 (1990) 1722-1725.

518) [\$K_s^0\$ production in \$\bar{p}p\$ interactions at \$\sqrt{s} = 630\$ GeV and 1800 GeV.](#)

By CDF Collaboration (F. Abe et al.).

[10.1103/PhysRevD.40.3791.](#)

Phys.Rev. D40 (1989) 3791-3794.

519) [Small Angle Physics At Cdf: A Progress Report.](#)

By CDF Collaboration (Riccardo Paoletti for the collaboration).

520) [Search for a light Higgs boson at the Tevatron \$\bar{p}p\$ collider.](#)

By CDF Collaboration (F. Abe et al.).

[10.1103/PhysRevD.41.1717.](#)

Phys.Rev. D41 (1990) 1717.

521) [Measurement of the ratio \$\sigma\(W \rightarrow e \nu\) / \sigma\(Z \rightarrow e e\)\$ in \$\bar{p}p\$ collisions at \$\sqrt{s} = 1.8\$ TeV.](#)

By CDF Collaboration (F. Abe et al.).

[10.1103/PhysRevLett.64.152.](#)

Phys.Rev.Lett. 64 (1990) 152-156.

522) [Two jet differential cross-section in \$\bar{p}p\$ collisions at \$\sqrt{s} = 1.8\$ TeV.](#)

By CDF Collaboration (F. Abe et al.).

[10.1103/PhysRevLett.64.157.](#)

Phys.Rev.Lett. 64 (1990) 157.

523) [Pseudorapidity distributions of charged particles produced in \$\bar{p}p\$ interactions at \$\sqrt{s} = 630\$ GeV and 1800 GeV.](#)

By CDF Collaboration (F. Abe et al.).

[10.1103/PhysRevD.41.2330.](#)

Phys.Rev. D41 (1990) 2330.

524) [A measurement of \$D^*\$ production in jets from \$\bar{p}p\$ collisions at \$\sqrt{s} = 1.8\$ TeV.](#)

By CDF Collaboration (F. Abe et al.).

[10.1103/PhysRevLett.64.348.](#)

Phys.Rev.Lett. 64 (1990) 348.

525) [A search for the top quark in the reaction \$\bar{p}p \rightarrow e + \text{jets}\$ at \$\sqrt{s} = 1.8\$ TeV.](#)

By CDF Collaboration (F. Abe et al.).

[10.1103/PhysRevLett.64.142.](#)

Phys.Rev.Lett. 64 (1990) 142-146.

526) [Measurement of the mass and width of the \$Z^0\$ boson at the Fermilab Tevatron.](#)

By CDF Collaboration (F. Abe et al.).

[10.1103/PhysRevLett.63.720.](#)

Phys.Rev.Lett. 63 (1989) 720.

- 527) [Search for Heavy Stable Particles at the Fermilab Collider.](#)
By CDF Collaboration (F. Abe et al.).
[10.1103/PhysRevLett.63.1447.](#)
Phys.Rev.Lett. 63 (1989) 1447.
- 528) [Dijet angular distributions from \$\bar{p}p\$ collisions at \$\sqrt{s} = 1.8\$ TeV.](#)
By CDF Collaboration (F. Abe et al.).
[10.1103/PhysRevLett.62.3020.](#)
Phys.Rev.Lett. 62 (1989) 3020-3023.
- 529) [Limits on the masses of supersymmetric particles from 1.8 TeV \$p\bar{p}\$ collisions.](#)
By CDF Collaboration (F. Abe et al.).
[10.1103/PhysRevLett.62.1825.](#)
Phys.Rev.Lett. 62 (1989) 1825-1828.
- 530) [Silicon Detectors For P Anti-p Small Angle Physics.](#)
By G. Tonelli, G. Apollinari, F. Bosi, S. Belforte, E. Focardi, R. Paoletti, F. Zetti.
[10.1016/0168-9002\(89\)90545-7.](#)
Nucl.Instrum.Meth. A277 (1989) 138-146.
- 531) [Measurement of the inclusive jet cross-section in \$\bar{p}p\$ collisions at \$\sqrt{s} = 1.8\$ TeV.](#)
By CDF Collaboration (F. Abe et al.).
[10.1103/PhysRevLett.62.613.](#)
Phys.Rev.Lett. 62 (1989) 613.
- 532) [A measurement of \$W\$ boson production in 1.8 TeV \$\bar{p}p\$ collisions.](#)
By CDF Collaboration (F. Abe et al.).
[10.1103/PhysRevLett.62.1005.](#)
Phys.Rev.Lett. 62 (1989) 1005.
- 533) [SILICON DETECTORS FOR \$p\bar{p}\$ ANGLE PHYSICS.](#)
By G. Apollinari, F. Bosi, S. Belforte, E. Focardi, R. Paoletti, G. Tonelli, F. Zetti.
- 534) [Transverse momentum distributions of charged particles produced in \$\bar{p}p\$ interactions at \$\sqrt{s} = 630\$ GeV and 1800 GeV.](#)
By CDF Collaboration (F. Abe et al.).
[10.1103/PhysRevLett.61.1819.](#)
Phys.Rev.Lett. 61 (1988) 1819.
- 535) [The CDF Detector: An Overview.](#)
By CDF Collaboration (F. Abe et al.).
[10.1016/0168-9002\(88\)90298-7.](#)
Nucl.Instrum.Meth. A271 (1988) 387-403.
- 536) [The CDF Central and Endwall Hadron Calorimeter.](#)
By CDF Collaboration (S. Bertolucci et al.).
[10.1016/0168-9002\(88\)90476-7.](#)
Nucl.Instrum.Meth. A267 (1988) 301.
- 537) [Proposal for upgrading the Collider Detector at Fermilab.](#)
By CDF Collaboration (R. Diebold et al.).
- 538) [The Cdf Leading Particle Spectrometers.](#)
By CDF Collaboration (Konstantin A. Goulios et al.).